

表 496 専従専門医の有無 体重(kg)別(透析患者全体)

体重	なし	あり	合計	記載なし	総計
20未満	18	25	43		43
(%)	(41.9)	(58.1)	(100.0)		
20～	229	273	502	1	503
(%)	(45.6)	(54.4)	(100.0)		
30～	7,434	8,409	15,843		15,843
(%)	(46.9)	(53.1)	(100.0)		
40～	28,649	35,224	63,873		63,873
(%)	(44.9)	(55.1)	(100.0)		
50～	38,518	50,547	89,065		89,065
(%)	(43.2)	(56.8)	(100.0)		
60～	26,874	37,928	64,802		64,802
(%)	(41.5)	(58.5)	(100.0)		
70～	11,641	17,042	28,683		28,683
(%)	(40.6)	(59.4)	(100.0)		
80～	4,184	6,469	10,653		10,653
(%)	(39.3)	(60.7)	(100.0)		
90～	1,556	2,462	4,018		4,018
(%)	(38.7)	(61.3)	(100.0)		
100～	549	804	1,353		1,353
(%)	(40.6)	(59.4)	(100.0)		
110～	225	376	601		601
(%)	(37.4)	(62.6)	(100.0)		
120～	169	269	438		438
(%)	(38.6)	(61.4)	(100.0)		
合計	120,046	159,828	279,874	1	279,875
(%)	(42.9)	(57.1)	(100.0)		
記載なし	13,294	16,939	30,233		30,233
(%)	(44.0)	(56.0)	(100.0)		
総計	133,340	176,767	310,107	1	310,108
(%)	(43.0)	(57.0)	(100.0)		
平均	57.47	58.51	58.06	21.30	58.06
標準偏差	13.31	13.57	13.47		13.47

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査