

表 497 専従専門医の有無 透析前BUN(mg/dL)別(透析患者全体)

透析前BUN	なし	あり	合計	記載なし	総計
20未満	426	403	829		829
(%)	(51.4)	(48.6)	(100.0)		
20～	2,095	2,399	4,494		4,494
(%)	(46.6)	(53.4)	(100.0)		
30～	8,027	9,627	17,654		17,654
(%)	(45.5)	(54.5)	(100.0)		
40～	19,422	24,505	43,927	1	43,928
(%)	(44.2)	(55.8)	(100.0)		
50～	29,315	39,321	68,636		68,636
(%)	(42.7)	(57.3)	(100.0)		
60～	28,433	39,520	67,953		67,953
(%)	(41.8)	(58.2)	(100.0)		
70～	18,896	25,960	44,856		44,856
(%)	(42.1)	(57.9)	(100.0)		
80～	8,970	12,045	21,015		21,015
(%)	(42.7)	(57.3)	(100.0)		
90～	3,342	4,426	7,768		7,768
(%)	(43.0)	(57.0)	(100.0)		
100～	1,016	1,301	2,317		2,317
(%)	(43.8)	(56.2)	(100.0)		
110～	309	386	695		695
(%)	(44.5)	(55.5)	(100.0)		
120～	188	207	395		395
(%)	(47.6)	(52.4)	(100.0)		
合計	120,439	160,100	280,539	1	280,540
(%)	(42.9)	(57.1)	(100.0)		
記載なし	12,901	16,667	29,568		29,568
(%)	(43.6)	(56.4)	(100.0)		
総計	133,340	176,767	310,107	1	310,108
(%)	(43.0)	(57.0)	(100.0)		
平均	60.46	60.92	60.72	45.00	60.72
標準偏差	16.14	15.69	15.89		15.89

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査