

表 524 専従専門医の有無 PTH値 (pg/mL) 別 (施設血液透析患者、PTH測定法:whole-PTH)

PTH値	なし	あり	合計	記載なし	総計
20未満	470	1,091	1,561		1,561
(%)	(30.1)	(69.9)	(100.0)		
20～	543	1,401	1,944		1,944
(%)	(27.9)	(72.1)	(100.0)		
35～	446	1,232	1,678		1,678
(%)	(26.6)	(73.4)	(100.0)		
50～	649	1,936	2,585		2,585
(%)	(25.1)	(74.9)	(100.0)		
75～	549	1,704	2,253		2,253
(%)	(24.4)	(75.6)	(100.0)		
100～	439	1,426	1,865		1,865
(%)	(23.5)	(76.5)	(100.0)		
125～	302	939	1,241		1,241
(%)	(24.3)	(75.7)	(100.0)		
150～	10	31	41		41
(%)	(24.4)	(75.6)	(100.0)		
151～	248	743	991		991
(%)	(25.0)	(75.0)	(100.0)		
180～	176	517	693		693
(%)	(25.4)	(74.6)	(100.0)		
210～	138	355	493		493
(%)	(28.0)	(72.0)	(100.0)		
240～	161	391	552		552
(%)	(29.2)	(70.8)	(100.0)		
300～	68	148	216		216
(%)	(31.5)	(68.5)	(100.0)		
350～	44	97	141		141
(%)	(31.2)	(68.8)	(100.0)		
400～	54	105	159		159
(%)	(34.0)	(66.0)	(100.0)		
500～	57	129	186		186
(%)	(30.6)	(69.4)	(100.0)		
合計	4,354	12,245	16,599		16,599
(%)	(26.2)	(73.8)	(100.0)		
記載なし	92	390	482		482
(%)	(19.1)	(80.9)	(100.0)		
総計	4,446	12,635	17,081		17,081
(%)	(26.0)	(74.0)	(100.0)		
平均	107.85	105.95	106.45		106.45
標準偏差	112.87	113.98	113.69		113.69

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査