

表 530 専従専門医の有無 透析前脈拍(拍/分)別(施設血液透析患者)

| 透析前脈拍 | なし | あり | 合計 | 記載なし | 総計 |
|-------|---------|---------|---------|------|---------|
| 40未満 | 76 | 84 | 160 | | 160 |
| (%) | (47.5) | (52.5) | (100.0) | | |
| 40～ | 982 | 1,055 | 2,037 | | 2,037 |
| (%) | (48.2) | (51.8) | (100.0) | | |
| 50～ | 7,477 | 9,021 | 16,498 | | 16,498 |
| (%) | (45.3) | (54.7) | (100.0) | | |
| 60～ | 22,552 | 28,254 | 50,806 | | 50,806 |
| (%) | (44.4) | (55.6) | (100.0) | | |
| 70～ | 26,281 | 34,346 | 60,627 | | 60,627 |
| (%) | (43.3) | (56.7) | (100.0) | | |
| 80～ | 16,429 | 21,760 | 38,189 | | 38,189 |
| (%) | (43.0) | (57.0) | (100.0) | | |
| 90～ | 7,064 | 9,505 | 16,569 | | 16,569 |
| (%) | (42.6) | (57.4) | (100.0) | | |
| 100～ | 2,139 | 2,934 | 5,073 | | 5,073 |
| (%) | (42.2) | (57.8) | (100.0) | | |
| 110～ | 477 | 721 | 1,198 | | 1,198 |
| (%) | (39.8) | (60.2) | (100.0) | | |
| 120～ | 191 | 323 | 514 | | 514 |
| (%) | (37.2) | (62.8) | (100.0) | | |
| 合計 | 83,668 | 108,003 | 191,671 | | 191,671 |
| (%) | (43.7) | (56.3) | (100.0) | | |
| 記載なし | 26,824 | 37,146 | 63,970 | | 63,970 |
| (%) | (41.9) | (58.1) | (100.0) | | |
| 総計 | 110,492 | 145,149 | 255,641 | | 255,641 |
| (%) | (43.2) | (56.8) | (100.0) | | |
| 平均 | 74.17 | 74.71 | 74.47 | | 74.47 |
| 標準偏差 | 12.74 | 12.82 | 12.79 | | 12.79 |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査