

表 2064 HDF希釈方法 透析前BUN(mg/dL)別（オフラインHDF 女性のみ 透析回数週3回 透析歴2年以上）

透析前BUN	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
20未満	1	4			5		2	7
(%)	(20.0)	(80.0)			(100.0)			
20～	4	18			22		2	24
(%)	(18.2)	(81.8)			(100.0)			
30～	13	83			96		14	110
(%)	(13.5)	(86.5)			(100.0)			
40～	24	234			258		61	319
(%)	(9.3)	(90.7)			(100.0)			
50～	53	380		1	434		78	512
(%)	(12.2)	(87.6)		(0.2)	(100.0)			
60～	56	397	1		454		65	519
(%)	(12.3)	(87.4)	(0.2)		(100.0)			
70～	38	289			327		69	396
(%)	(11.6)	(88.4)			(100.0)			
80～	14	120			134		38	172
(%)	(10.4)	(89.6)			(100.0)			
90～	4	41	1		46		8	54
(%)	(8.7)	(89.1)	(2.2)		(100.0)			
100～	2	21			23			23
(%)	(8.7)	(91.3)			(100.0)			
110～	1				1		2	3
(%)	(100.0)				(100.0)			
120～		2			2			2
(%)		(100.0)			(100.0)			
合計	210	1,589	2	1	1,802		339	2,141
(%)	(11.7)	(88.2)	(0.1)	(0.1)	(100.0)			
記載なし	4	22			26		18	44
(%)	(15.4)	(84.6)			(100.0)			
総計	214	1,611	2	1	1,828		357	2,185
(%)	(11.7)	(88.1)	(0.1)	(0.1)	(100.0)			
平均	61.81	61.75	76.50	58.00	61.77		62.22	61.84
標準偏差	15.84	15.57	23.33		15.60		15.90	15.64

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査