

表 589 糖尿病の既往 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	50,767	6,854	57,621	69	10,887	68,577
(%)	(88.1)	(11.9)	(100.0)			
慢性腎盂腎炎	1,597	252	1,849	4	312	2,165
(%)	(86.4)	(13.6)	(100.0)			
急速進行性糸球体腎炎	1,204	371	1,575	3	256	1,834
(%)	(76.4)	(23.6)	(100.0)			
妊娠腎/妊娠中毒症	821	83	904	4	156	1,064
(%)	(90.8)	(9.2)	(100.0)			
その他の分類不能の腎炎	684	108	792		157	949
(%)	(86.4)	(13.6)	(100.0)			
多発性嚢胞腎	6,386	554	6,940	12	1,246	8,198
(%)	(92.0)	(8.0)	(100.0)			
腎硬化症	15,542	3,006	18,548	51	2,861	21,460
(%)	(83.8)	(16.2)	(100.0)			
悪性高血圧	1,394	192	1,586	2	293	1,881
(%)	(87.9)	(12.1)	(100.0)			
糖尿病性腎症	39	89,314	89,353		1,154	90,507
(%)	(0.0)	(100.0)	(100.0)			
SLE腎炎	1,114	213	1,327	2	221	1,550
(%)	(83.9)	(16.1)	(100.0)			
アミロイド腎	221	53	274	3	48	325
(%)	(80.7)	(19.3)	(100.0)			
痛風腎	638	82	720	1	107	828
(%)	(88.6)	(11.4)	(100.0)			
先天性代謝異常による腎不全	135	19	154		23	177
(%)	(87.7)	(12.3)	(100.0)			
腎・尿路結核	118	11	129		29	158
(%)	(91.5)	(8.5)	(100.0)			
腎・尿路結石	316	44	360		70	430
(%)	(87.8)	(12.2)	(100.0)			
腎・尿路腫瘍	490	81	571		115	686
(%)	(85.8)	(14.2)	(100.0)			
閉塞性尿路障害	413	39	452	1	64	517
(%)	(91.4)	(8.6)	(100.0)			
骨髄腫	149	20	169	1	20	190
(%)	(88.2)	(11.8)	(100.0)			
腎形成不全	341	24	365		54	419
(%)	(93.4)	(6.6)	(100.0)			
不明	13,470	3,253	16,723	54	3,348	20,125
(%)	(80.5)	(19.5)	(100.0)			
再導入	1,019	211	1,230	1	186	1,417
(%)	(82.8)	(17.2)	(100.0)			
その他	3,600	1,092	4,692	22	800	5,514
(%)	(76.7)	(23.3)	(100.0)			
合計	100,458	105,876	206,334	230	22,407	228,971
(%)	(48.7)	(51.3)	(100.0)			
記載なし						
(%)						
総計	100,458	105,876	206,334	230	22,407	228,971
(%)	(48.7)	(51.3)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査