

表 602 糖尿病の既往 身長(cm)別（施設血液透析患者 透析回数週3回）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	105	117	222		15	237
(%)	(47.3)	(52.7)	(100.0)			
130～	326	192	518		50	568
(%)	(62.9)	(37.1)	(100.0)			
135～	1,254	824	2,078		227	2,305
(%)	(60.3)	(39.7)	(100.0)			
140～	4,369	3,177	7,546	13	724	8,283
(%)	(57.9)	(42.1)	(100.0)			
145～	9,591	7,489	17,080	20	1,719	18,819
(%)	(56.2)	(43.8)	(100.0)			
150～	14,760	12,431	27,191	32	2,791	30,014
(%)	(54.3)	(45.7)	(100.0)			
155～	16,505	15,292	31,797	28	3,032	34,857
(%)	(51.9)	(48.1)	(100.0)			
160～	18,212	20,611	38,823	28	3,407	42,258
(%)	(46.9)	(53.1)	(100.0)			
165～	15,622	19,071	34,693	20	2,863	37,576
(%)	(45.0)	(55.0)	(100.0)			
170～	9,799	12,147	21,946	19	1,862	23,827
(%)	(44.7)	(55.3)	(100.0)			
175～	3,389	4,492	7,881	5	663	8,549
(%)	(43.0)	(57.0)	(100.0)			
180～	854	1,233	2,087	2	165	2,254
(%)	(40.9)	(59.1)	(100.0)			
185～	132	194	326		25	351
(%)	(40.5)	(59.5)	(100.0)			
190～	14	40	54		3	57
(%)	(25.9)	(74.1)	(100.0)			
合計	94,932	97,310	192,242	167	17,546	209,955
(%)	(49.4)	(50.6)	(100.0)			
記載なし	5,526	8,566	14,092	63	4,861	19,016
(%)	(39.2)	(60.8)	(100.0)			
総計	100,458	105,876	206,334	230	22,407	228,971
(%)	(48.7)	(51.3)	(100.0)			
平均	159.12	160.87	160.00	158.23	159.31	159.94
標準偏差	9.63	9.47	9.59	9.68	9.60	9.59

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査