

表 703 心筋梗塞の既往 原疾患別（透析患者全体）

| 原疾患           | 既往なし    | 既往あり   | 合計      | 不明  | 記載なし   | 総計      |
|---------------|---------|--------|---------|-----|--------|---------|
| 慢性糸球体腎炎       | 68,057  | 5,540  | 73,597  | 226 | 23,147 | 96,970  |
| (%)           | (92.5)  | (7.5)  | (100.0) |     |        |         |
| 慢性腎盂腎炎        | 2,207   | 152    | 2,359   | 5   | 678    | 3,042   |
| (%)           | (93.6)  | (6.4)  | (100.0) |     |        |         |
| 急速進行性糸球体腎炎    | 1,786   | 115    | 1,901   | 7   | 586    | 2,494   |
| (%)           | (94.0)  | (6.0)  | (100.0) |     |        |         |
| 妊娠腎/妊娠中毒症     | 1,202   | 64     | 1,266   | 6   | 342    | 1,614   |
| (%)           | (94.9)  | (5.1)  | (100.0) |     |        |         |
| その他の分類不能の腎炎   | 975     | 52     | 1,027   | 3   | 343    | 1,373   |
| (%)           | (94.9)  | (5.1)  | (100.0) |     |        |         |
| 多発性嚢胞腎        | 7,941   | 524    | 8,465   | 24  | 2,517  | 11,006  |
| (%)           | (93.8)  | (6.2)  | (100.0) |     |        |         |
| 腎硬化症          | 19,788  | 2,354  | 22,142  | 119 | 6,037  | 28,298  |
| (%)           | (89.4)  | (10.6) | (100.0) |     |        |         |
| 悪性高血圧         | 1,795   | 158    | 1,953   | 6   | 614    | 2,573   |
| (%)           | (91.9)  | (8.1)  | (100.0) |     |        |         |
| 糖尿病性腎症        | 77,626  | 12,645 | 90,271  | 380 | 27,430 | 118,081 |
| (%)           | (86.0)  | (14.0) | (100.0) |     |        |         |
| SLE腎炎         | 1,636   | 103    | 1,739   | 7   | 490    | 2,236   |
| (%)           | (94.1)  | (5.9)  | (100.0) |     |        |         |
| アミロイド腎        | 313     | 16     | 329     | 3   | 119    | 451     |
| (%)           | (95.1)  | (4.9)  | (100.0) |     |        |         |
| 痛風腎           | 821     | 62     | 883     | 4   | 226    | 1,113   |
| (%)           | (93.0)  | (7.0)  | (100.0) |     |        |         |
| 先天性代謝異常による腎不全 | 196     | 8      | 204     | 1   | 67     | 272     |
| (%)           | (96.1)  | (3.9)  | (100.0) |     |        |         |
| 腎・尿路結核        | 141     | 12     | 153     |     | 53     | 206     |
| (%)           | (92.2)  | (7.8)  | (100.0) |     |        |         |
| 腎・尿路結石        | 408     | 35     | 443     |     | 138    | 581     |
| (%)           | (92.1)  | (7.9)  | (100.0) |     |        |         |
| 腎・尿路腫瘍        | 618     | 38     | 656     | 2   | 228    | 886     |
| (%)           | (94.2)  | (5.8)  | (100.0) |     |        |         |
| 閉塞性尿路障害       | 534     | 34     | 568     | 3   | 152    | 723     |
| (%)           | (94.0)  | (6.0)  | (100.0) |     |        |         |
| 骨髄腫           | 197     | 16     | 213     | 1   | 62     | 276     |
| (%)           | (92.5)  | (7.5)  | (100.0) |     |        |         |
| 腎形成不全         | 496     | 13     | 509     |     | 140    | 649     |
| (%)           | (97.4)  | (2.6)  | (100.0) |     |        |         |
| 不明            | 18,596  | 1,684  | 20,280  | 82  | 7,182  | 27,544  |
| (%)           | (91.7)  | (8.3)  | (100.0) |     |        |         |
| 再導入           | 1,533   | 120    | 1,653   | 1   | 509    | 2,163   |
| (%)           | (92.7)  | (7.3)  | (100.0) |     |        |         |
| その他           | 5,222   | 451    | 5,673   | 37  | 1,847  | 7,557   |
| (%)           | (92.1)  | (7.9)  | (100.0) |     |        |         |
| 合計            | 212,088 | 24,196 | 236,284 | 917 | 72,907 | 310,108 |
| (%)           | (89.8)  | (10.2) | (100.0) |     |        |         |
| 記載なし          |         |        |         |     |        |         |
| (%)           |         |        |         |     |        |         |
| 総計            | 212,088 | 24,196 | 236,284 | 917 | 72,907 | 310,108 |
| (%)           | (89.8)  | (10.2) | (100.0) |     |        |         |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査