

表 1001 脳梗塞の既往 PTH値(pg/mL)別（透析患者全体）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	1,301	322	1,623	3	268	1,894
(%)	(80.2)	(19.8)	(100.0)			
20～	1,607	400	2,007	1	323	2,331
(%)	(80.1)	(19.9)	(100.0)			
35～	1,439	327	1,766	3	274	2,043
(%)	(81.5)	(18.5)	(100.0)			
50～	2,174	506	2,680	3	470	3,153
(%)	(81.1)	(18.9)	(100.0)			
75～	1,877	457	2,334	3	411	2,748
(%)	(80.4)	(19.6)	(100.0)			
100～	1,552	366	1,918	1	379	2,298
(%)	(80.9)	(19.1)	(100.0)			
125～	1,002	257	1,259		280	1,539
(%)	(79.6)	(20.4)	(100.0)			
150～	34	3	37		11	48
(%)	(91.9)	(8.1)	(100.0)			
151～	879	187	1,066		197	1,263
(%)	(82.5)	(17.5)	(100.0)			
180～	606	137	743		132	875
(%)	(81.6)	(18.4)	(100.0)			
210～	437	94	531		97	628
(%)	(82.3)	(17.7)	(100.0)			
240～	468	98	566		108	674
(%)	(82.7)	(17.3)	(100.0)			
300～	196	35	231		53	284
(%)	(84.8)	(15.2)	(100.0)			
350～	129	24	153		26	179
(%)	(84.3)	(15.7)	(100.0)			
400～	147	23	170		32	202
(%)	(86.5)	(13.5)	(100.0)			
500～	172	40	212	1	34	247
(%)	(81.1)	(18.9)	(100.0)			
合計	14,020	3,276	17,296	15	3,095	20,406
(%)	(81.1)	(18.9)	(100.0)			
記載なし	311	85	396	1	260	657
(%)	(78.5)	(21.5)	(100.0)			
総計	14,331	3,361	17,692	16	3,355	21,063
(%)	(81.0)	(19.0)	(100.0)			
平均	108.19	103.73	107.35	93.13	111.76	108.01
標準偏差	114.30	110.11	113.53	167.40	116.38	114.01

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査