

表 1159 四肢切断の有無 透析前BUN(mg/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前BUN	切断なし	切断あり	合計	不明	記載なし	総計
20未満	368	31	399	1	79	479
(%)	(92.2)	(7.8)	(100.0)			
20～	1,934	132	2,066	3	411	2,480
(%)	(93.6)	(6.4)	(100.0)			
30～	8,060	562	8,622	15	1,721	10,358
(%)	(93.5)	(6.5)	(100.0)			
40～	21,317	1,133	22,450	26	4,335	26,811
(%)	(95.0)	(5.0)	(100.0)			
50～	34,068	1,480	35,548	31	7,080	42,659
(%)	(95.8)	(4.2)	(100.0)			
60～	34,218	1,402	35,620	34	6,989	42,643
(%)	(96.1)	(3.9)	(100.0)			
70～	22,974	751	23,725	30	4,567	28,322
(%)	(96.8)	(3.2)	(100.0)			
80～	10,731	338	11,069	14	2,129	13,212
(%)	(96.9)	(3.1)	(100.0)			
90～	3,870	130	4,000	7	796	4,803
(%)	(96.8)	(3.3)	(100.0)			
100～	1,111	39	1,150	3	195	1,348
(%)	(96.6)	(3.4)	(100.0)			
110～	320	14	334		60	394
(%)	(95.8)	(4.2)	(100.0)			
120～	163	9	172		32	204
(%)	(94.8)	(5.2)	(100.0)			
合計	139,134	6,021	145,155	164	28,394	173,713
(%)	(95.9)	(4.1)	(100.0)			
記載なし	1,805	149	1,954	5	2,537	4,496
(%)	(92.4)	(7.6)	(100.0)			
総計	140,939	6,170	147,109	169	30,931	178,209
(%)	(95.8)	(4.2)	(100.0)			
平均	61.10	57.82	60.97	61.30	60.81	60.94
標準偏差	15.62	16.02	15.65	17.05	15.56	15.64

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査