

表 1373 大腿骨頸部骨折の既往 PTH値(pg/mL)別（血液透析濾過患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	207	8	215		34	249
(%)	(96.3)	(3.7)	(100.0)			
20～	254	8	262		39	301
(%)	(96.9)	(3.1)	(100.0)			
35～	226	10	236		38	274
(%)	(95.8)	(4.2)	(100.0)			
50～	358	12	370		74	444
(%)	(96.8)	(3.2)	(100.0)			
75～	311	10	321		70	391
(%)	(96.9)	(3.1)	(100.0)			
100～	269	8	277		54	331
(%)	(97.1)	(2.9)	(100.0)			
125～	187	7	194		40	234
(%)	(96.4)	(3.6)	(100.0)			
150～	4		4			4
(%)	(100.0)		(100.0)			
151～	178	6	184		31	215
(%)	(96.7)	(3.3)	(100.0)			
180～	116	3	119		18	137
(%)	(97.5)	(2.5)	(100.0)			
210～	79	6	85		13	98
(%)	(92.9)	(7.1)	(100.0)			
240～	74	5	79		9	88
(%)	(93.7)	(6.3)	(100.0)			
300～	36	1	37		5	42
(%)	(97.3)	(2.7)	(100.0)			
350～	27		27		2	29
(%)	(100.0)		(100.0)			
400～	19	1	20		4	24
(%)	(95.0)	(5.0)	(100.0)			
500～	41	1	42		3	45
(%)	(97.6)	(2.4)	(100.0)			
合計	2,386	86	2,472		434	2,906
(%)	(96.5)	(3.5)	(100.0)			
記載なし	7	1	8		23	31
(%)	(87.5)	(12.5)	(100.0)			
総計	2,393	87	2,480		457	2,937
(%)	(96.5)	(3.5)	(100.0)			
平均	113.05	113.60	113.07		104.49	111.79
標準偏差	111.51	96.97	111.02		89.55	108.12

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査