

表 1814 腎移植の回数 透析後体重(kg)別（施設血液透析患者 透析回数週3回）

透析後体重	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
20未満	67					67	1	16	84
(%)	(100.0)					(100.0)			
20～	642	8	1			651	3	148	802
(%)	(98.6)	(1.2)	(0.2)			(100.0)			
30～	15,333	141	10	1		15,485	74	3,558	19,117
(%)	(99.0)	(0.9)	(0.1)	(0.0)		(100.0)			
40～	47,673	535	29	4		48,241	185	11,383	59,809
(%)	(98.8)	(1.1)	(0.1)	(0.0)		(100.0)			
50～	57,277	601	24			57,902	232	13,975	72,109
(%)	(98.9)	(1.0)	(0.0)			(100.0)			
60～	35,070	306	18			35,394	109	8,579	44,082
(%)	(99.1)	(0.9)	(0.1)			(100.0)			
70～	13,681	89	3			13,773	40	3,371	17,184
(%)	(99.3)	(0.6)	(0.0)			(100.0)			
80～	4,818	37				4,855	12	1,214	6,081
(%)	(99.2)	(0.8)				(100.0)			
90～	1,735	15				1,750	5	440	2,195
(%)	(99.1)	(0.9)				(100.0)			
100～	591	4	1			596	2	157	755
(%)	(99.2)	(0.7)	(0.2)			(100.0)			
110～	227	2				229	2	68	299
(%)	(99.1)	(0.9)				(100.0)			
120～	151					151		38	189
(%)	(100.0)					(100.0)			
合計	177,265	1,738	86	5		179,094	665	42,947	222,706
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
記載なし	3,506	26	2			3,534	22	2,709	6,265
(%)	(99.2)	(0.7)	(0.1)			(100.0)			
総計	180,771	1,764	88	5		182,628	687	45,656	228,971
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
平均	55.41	54.04	52.33	42.40		55.40	53.67	55.67	55.45
標準偏差	12.90	11.75	12.12	4.73		12.89	12.41	12.95	12.90

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査