

表 1825 腎移植の回数 透析前CRP濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

透析前CRP濃度	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
0.2未満	85,478	868	49	1		86,396	309	18,824	105,529
(%)	(98.9)	(1.0)	(0.1)	(0.0)		(100.0)			
0.2～	28,878	271	8	2		29,159	101	6,118	35,378
(%)	(99.0)	(0.9)	(0.0)	(0.0)		(100.0)			
0.5～	14,614	127	8			14,749	60	3,338	18,147
(%)	(99.1)	(0.9)	(0.1)			(100.0)			
1.0～	9,606	91	5			9,702	34	2,036	11,772
(%)	(99.0)	(0.9)	(0.1)			(100.0)			
2.0～	5,829	56	5	1		5,891	28	1,335	7,254
(%)	(98.9)	(1.0)	(0.1)	(0.0)		(100.0)			
4.0～	2,078	19	1			2,098	15	469	2,582
(%)	(99.0)	(0.9)	(0.0)			(100.0)			
6.0～	1,039	4				1,043	5	234	1,282
(%)	(99.6)	(0.4)				(100.0)			
8.0～	563	2				565	4	125	694
(%)	(99.6)	(0.4)				(100.0)			
10.0～	679	9	1			689	6	147	842
(%)	(98.5)	(1.3)	(0.1)			(100.0)			
15.0～	434	3				437	3	120	560
(%)	(99.3)	(0.7)				(100.0)			
合計	149,198	1,450	77	4		150,729	565	32,746	184,040
(%)	(99.0)	(1.0)	(0.1)	(0.0)		(100.0)			
記載なし	31,573	314	11	1		31,899	122	12,910	44,931
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
総計	180,771	1,764	88	5		182,628	687	45,656	228,971
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
平均	0.64	0.57	0.60	0.95		0.64	0.90	0.66	0.64
標準偏差	1.91	1.55	1.54	1.44		1.91	2.39	2.09	1.94

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査