

表 1876 腎移植の回数 BUN(mg/dL)別（腹膜透析患者）

BUN	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
20未満	22					22		7	29
(%)	(100.0)					(100.0)			
20～	164	2				166		17	183
(%)	(98.8)	(1.2)				(100.0)			
30～	575	3				578	1	93	672
(%)	(99.5)	(0.5)				(100.0)			
40～	1,195	13	1			1,209	2	181	1,392
(%)	(98.8)	(1.1)	(0.1)			(100.0)			
50～	1,409	6	1			1,416	5	192	1,613
(%)	(99.5)	(0.4)	(0.1)			(100.0)			
60～	988	13				1,001	1	148	1,150
(%)	(98.7)	(1.3)				(100.0)			
70～	479	4				483	2	64	549
(%)	(99.2)	(0.8)				(100.0)			
80～	219	5				224		21	245
(%)	(97.8)	(2.2)				(100.0)			
90～	79	1				80		9	89
(%)	(98.8)	(1.3)				(100.0)			
100～	31					31		9	40
(%)	(100.0)					(100.0)			
110～	6					6		1	7
(%)	(100.0)					(100.0)			
120～	6					6		1	7
(%)	(100.0)					(100.0)			
合計	5,173	47	2			5,222	11	743	5,976
(%)	(99.1)	(0.9)	(0.0)			(100.0)			
記載なし	602	9		1		612	12	2,341	2,965
(%)	(98.4)	(1.5)		(0.2)		(100.0)			
総計	5,775	56	2	1		5,834	23	3,084	8,941
(%)	(99.0)	(1.0)	(0.0)	(0.0)		(100.0)			
平均	54.96	56.68	45.50			54.97	53.91	54.45	54.90
標準偏差	15.66	16.74	6.36			15.67	12.45	16.91	15.82

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査