

表 1958 血流量(ml/分) 身長(cm)別 (施設血液透析患者 透析回数週3回)

身長	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	280～	300～	350～	400～	合計	記載なし	総計	平均	標準偏差	
130未満	1	2	7	56	13	45	83	12	10	1	1				231	6	237	181.73	31.49	
(%)	(0.4)	(0.9)	(3.0)	(24.2)	(5.6)	(19.5)	(35.9)	(5.2)	(4.3)	(0.4)	(0.4)				(100.0)					
130～		7	24	156	44	117	184	14	14			1	1		562	6	568	176.65	29.55	
(%)		(1.2)	(4.3)	(27.8)	(7.8)	(20.8)	(32.7)	(2.5)	(2.5)			(0.2)	(0.2)		(100.0)					
135～	2	22	70	532	154	535	782	77	60	4	2		21	4	11	2,276	29	2,305	182.57	37.71
(%)	(0.1)	(1.0)	(3.1)	(23.4)	(6.8)	(23.5)	(34.4)	(3.4)	(2.6)	(0.2)	(0.1)		(0.9)	(0.2)	(0.5)	(100.0)				
140～	4	47	202	1,553	514	1,814	3,254	415	288	19	16		26	4	20	8,176	107	8,283	185.52	32.56
(%)	(0.0)	(0.6)	(2.5)	(19.0)	(6.3)	(22.2)	(39.8)	(5.1)	(3.5)	(0.2)	(0.2)		(0.3)	(0.0)	(0.2)	(100.0)				
145～	4	85	328	2,678	967	3,918	8,223	1,215	864	63	48		105	9	59	18,566	253	18,819	190.61	32.98
(%)	(0.0)	(0.5)	(1.8)	(14.4)	(5.2)	(21.1)	(44.3)	(6.5)	(4.7)	(0.3)	(0.3)		(0.6)	(0.0)	(0.3)	(100.0)				
150～	6	127	363	3,304	1,143	5,387	14,363	2,548	1,802	142	107		204	30	84	29,610	404	30,014	195.30	32.68
(%)	(0.0)	(0.4)	(1.2)	(11.2)	(3.9)	(18.2)	(48.5)	(8.6)	(6.1)	(0.5)	(0.4)		(0.7)	(0.1)	(0.3)	(100.0)				
155～	5	96	307	2,652	938	5,446	17,061	4,011	2,999	243	175		383	27	84	34,427	430	34,857	200.87	32.31
(%)	(0.0)	(0.3)	(0.9)	(7.7)	(2.7)	(15.8)	(49.6)	(11.7)	(8.7)	(0.7)	(0.5)		(1.1)	(0.1)	(0.2)	(100.0)				
160～	5	117	266	2,422	828	5,138	19,874	6,227	5,118	463	377		673	70	111	41,689	569	42,258	206.74	33.67
(%)	(0.0)	(0.3)	(0.6)	(5.8)	(2.0)	(12.3)	(47.7)	(14.9)	(12.3)	(1.1)	(0.9)		(1.6)	(0.2)	(0.3)	(100.0)				
165～	5	91	201	1,585	549	3,358	16,238	6,480	6,231	612	605		936	94	105	37,090	486	37,576	213.29	35.21
(%)	(0.0)	(0.2)	(0.5)	(4.3)	(1.5)	(9.1)	(43.8)	(17.5)	(16.8)	(1.7)	(1.6)		(2.5)	(0.3)	(0.3)	(100.0)				
170～		42	114	746	221	1,611	8,967	4,540	5,094	501	561		923	74	80	23,474	353	23,827	220.01	36.61
(%)		(0.2)	(0.5)	(3.2)	(0.9)	(6.9)	(38.2)	(19.3)	(21.7)	(2.1)	(2.4)		(3.9)	(0.3)	(0.3)	(100.0)				
175～	1	17	29	213	68	458	2,713	1,706	2,143	265	239		529	46	27	8,454	95	8,549	226.36	37.98
(%)	(0.0)	(0.2)	(0.3)	(2.5)	(0.8)	(5.4)	(32.1)	(20.2)	(25.3)	(3.1)	(2.8)		(6.3)	(0.5)	(0.3)	(100.0)				
180～		2	11	46	13	92	552	398	704	86	89		193	34	4	2,224	30	2,254	234.67	40.01
(%)		(0.1)	(0.5)	(2.1)	(0.6)	(4.1)	(24.8)	(17.9)	(31.7)	(3.9)	(4.0)		(8.7)	(1.5)	(0.2)	(100.0)				
185～		1		6	1	15	67	69	116	15	17		38	3		348	3	351	238.32	37.58
(%)		(0.3)		(1.7)	(0.3)	(4.3)	(19.3)	(19.8)	(33.3)	(4.3)	(4.9)		(10.9)	(0.9)		(100.0)				
190～			2	1	1	17	8	9	9	3	1		11	3		56	1	57	241.43	49.26
(%)			(3.6)	(1.8)	(1.8)	(30.4)	(14.3)	(16.1)	(5.4)	(1.8)	(1.8)		(19.6)	(5.4)		(100.0)				
合計	33	656	1,922	15,951	5,454	27,935	92,378	27,720	25,452	2,417	2,238		4,043	399	585	207,183	2,772	209,955	205.31	35.95
(%)	(0.0)	(0.3)	(0.9)	(7.7)	(2.6)	(13.5)	(44.6)	(13.4)	(12.3)	(1.2)	(1.1)		(2.0)	(0.2)	(0.3)	(100.0)				
記載なし	11	83	226	2,022	447	2,550	8,112	1,826	1,608	116	99		222	30	133	17,485	1,531	19,016	200.52	41.33
(%)	(0.1)	(0.5)	(1.3)	(11.6)	(2.6)	(14.6)	(46.4)	(10.4)	(9.2)	(0.7)	(0.6)		(1.3)	(0.2)	(0.8)	(100.0)				
総計	44	739	2,148	17,973	5,901	30,485	100,490	29,546	27,060	2,533	2,337		4,265	429	718	224,668	4,303	228,971	204.94	36.42
(%)	(0.0)	(0.3)	(1.0)	(8.0)	(2.6)	(13.6)	(44.7)	(13.2)	(12.0)	(1.1)	(1.0)		(1.9)	(0.2)	(0.3)	(100.0)				
平均	153.00	156.21	154.37	154.34	154.41	156.33	159.45	163.26	164.91	166.06	166.96		166.78	160.10	159.94	159.94	159.91	159.94		
標準偏差	11.58	10.28	10.32	9.72	9.22	9.14	8.94	8.50	8.60	8.61	8.12		8.81	9.94	9.56	9.59	9.60	9.59		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査