

表 2408 身長(cm) 総コレステロール濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

総コレステロール濃度	130未満	130～	135～	140～	145～	150～	155～	160～	165～	170～	175～	180～	185～	190～	合計	記載なし	総計	平均	標準偏差
100未満	2		4	16	56	92	124	225	192	145	59	18			933	59	992	162.64	8.72
(%)	(0.2)		(0.4)	(1.7)	(6.0)	(9.9)	(13.3)	(24.1)	(20.6)	(15.5)	(6.3)	(1.9)			(100.0)				
100～	1	3	13	45	125	269	445	607	651	433	167	47	6	1	2,813	175	2,988	162.99	8.65
(%)	(0.0)	(0.1)	(0.5)	(1.6)	(4.4)	(9.6)	(15.8)	(21.6)	(23.1)	(15.4)	(5.9)	(1.7)	(0.2)	(0.0)	(100.0)				
120～	3	7	47	124	338	546	791	1,158	1,118	793	344	76	10	2	5,357	337	5,694	162.23	9.36
(%)	(0.1)	(0.1)	(0.9)	(2.3)	(6.3)	(10.2)	(14.8)	(21.6)	(20.9)	(14.8)	(6.4)	(1.4)	(0.2)	(0.0)	(100.0)				
140～	11	16	44	186	502	905	996	1,340	1,239	858	339	85	13		6,534	371	6,905	161.00	9.72
(%)	(0.2)	(0.2)	(0.7)	(2.8)	(7.7)	(13.9)	(15.2)	(20.5)	(19.0)	(13.1)	(5.2)	(1.3)	(0.2)		(100.0)				
160～	5	12	49	224	507	841	939	1,055	998	741	260	66	13	2	5,712	299	6,011	160.29	9.62
(%)	(0.1)	(0.2)	(0.9)	(3.9)	(8.9)	(14.7)	(16.4)	(18.5)	(17.5)	(13.0)	(4.6)	(1.2)	(0.2)	(0.0)	(100.0)				
180～	4	8	33	131	372	633	669	634	555	404	156	52	8	1	3,660	213	3,873	159.50	9.63
(%)	(0.1)	(0.2)	(0.9)	(3.6)	(10.2)	(17.3)	(18.3)	(17.3)	(15.2)	(11.0)	(4.3)	(1.4)	(0.2)	(0.0)	(100.0)				
200～	3	4	22	79	232	436	405	376	295	172	73	28	5		2,130	123	2,253	158.46	9.34
(%)	(0.1)	(0.2)	(1.0)	(3.7)	(10.9)	(20.5)	(19.0)	(17.7)	(13.8)	(8.1)	(3.4)	(1.3)	(0.2)		(100.0)				
220～		2	9	40	110	194	185	146	103	64	33	2	2		890	66	956	157.55	8.99
(%)		(0.2)	(1.0)	(4.5)	(12.4)	(21.8)	(20.8)	(16.4)	(11.6)	(7.2)	(3.7)	(0.2)	(0.2)		(100.0)				
240～	1	1	3	19	41	82	80	50	43	30	9	4			363	21	384	157.53	9.25
(%)	(0.3)	(0.3)	(0.8)	(5.2)	(11.3)	(22.6)	(22.0)	(13.8)	(11.8)	(8.3)	(2.5)	(1.1)			(100.0)				
260～		1		10	33	41	43	23	21	16	3	2			193	23	216	156.76	8.91
(%)		(0.5)		(5.2)	(17.1)	(21.2)	(22.3)	(11.9)	(10.9)	(8.3)	(1.6)	(1.0)			(100.0)				
合計	30	54	224	874	2,316	4,039	4,677	5,614	5,215	3,656	1,443	380	57	6	28,585	1,687	30,272	160.78	9.54
(%)	(0.1)	(0.2)	(0.8)	(3.1)	(8.1)	(14.1)	(16.4)	(19.6)	(18.2)	(12.8)	(5.0)	(1.3)	(0.2)	(0.0)	(100.0)				
記載なし	8	16	65	289	679	1,173	1,402	1,563	1,507	1,129	343	113	20	3	8,310	1,364	9,674	160.60	9.51
(%)	(0.1)	(0.2)	(0.8)	(3.5)	(8.2)	(14.1)	(16.9)	(18.8)	(18.1)	(13.6)	(4.1)	(1.4)	(0.2)	(0.0)	(100.0)				
総計	38	70	289	1,163	2,995	5,212	6,079	7,177	6,722	4,785	1,786	493	77	9	36,895	3,051	39,946	160.74	9.54
(%)	(0.1)	(0.2)	(0.8)	(3.2)	(8.1)	(14.1)	(16.5)	(19.5)	(18.2)	(13.0)	(4.8)	(1.3)	(0.2)	(0.0)	(100.0)				
平均	160.70	166.28	162.54	167.28	167.25	166.74	161.58	154.31	152.26	152.14	151.18	153.09	158.98	151.33	157.98	158.46	158.00		
標準偏差	37.35	34.93	33.57	35.14	36.54	36.48	36.86	34.75	33.80	33.16	32.67	34.44	31.03	31.45	35.56	37.88	35.70		

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査