

表 2465 透析後体重(kg) 体重減少率(%)別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

体重減少率	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
1.0未満		37	452	1,113	1,172	638	210	55	7	8		4	3,696		3,696	52.66	12.05
(%)		(1.0)	(12.2)	(30.1)	(31.7)	(17.3)	(5.7)	(1.5)	(0.2)	(0.2)		(0.1)	(100.0)				
1.0～		25	613	1,890	2,154	1,305	436	124	42	12	8		6,609		6,609	54.40	12.00
(%)		(0.4)	(9.3)	(28.6)	(32.6)	(19.7)	(6.6)	(1.9)	(0.6)	(0.2)	(0.1)		(100.0)				
2.0～		59	1,305	4,281	5,594	3,611	1,353	476	146	51	21	24	16,921		16,921	55.99	12.75
(%)		(0.3)	(7.7)	(25.3)	(33.1)	(21.3)	(8.0)	(2.8)	(0.9)	(0.3)	(0.1)	(0.1)	(100.0)				
3.0～		83	2,219	8,186	11,277	7,719	3,287	1,147	431	135	60	40	34,584		34,584	57.20	13.08
(%)		(0.2)	(6.4)	(23.7)	(32.6)	(22.3)	(9.5)	(3.3)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
4.0～	1	115	3,116	11,592	15,057	10,173	4,007	1,464	574	197	72	47	46,415		46,415	56.76	13.00
(%)	(0.0)	(0.2)	(6.7)	(25.0)	(32.4)	(21.9)	(8.6)	(3.2)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
5.0～		102	3,184	10,419	12,146	6,661	2,516	820	280	101	40	20	36,289		36,289	54.67	12.32
(%)		(0.3)	(8.8)	(28.7)	(33.5)	(18.4)	(6.9)	(2.3)	(0.8)	(0.3)	(0.1)	(0.1)	(100.0)				
6.0～		95	2,376	6,118	5,798	2,734	901	270	97	25	10	3	18,427		18,427	52.09	11.67
(%)		(0.5)	(12.9)	(33.2)	(31.5)	(14.8)	(4.9)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)	(100.0)				
7.0～		64	1,290	2,594	2,012	801	238	72	27	8	2		7,108		7,108	49.75	11.18
(%)		(0.9)	(18.1)	(36.5)	(28.3)	(11.3)	(3.3)	(1.0)	(0.4)	(0.1)	(0.0)		(100.0)				
8.0～		36	504	835	536	204	61	8	7	2	2		2,195		2,195	47.80	10.91
(%)		(1.6)	(23.0)	(38.0)	(24.4)	(9.3)	(2.8)	(0.4)	(0.3)	(0.1)	(0.1)		(100.0)				
9.0～	1	28	285	448	309	102	24	6	1	1			1,205		1,205	47.11	10.68
(%)	(0.1)	(2.3)	(23.7)	(37.2)	(25.6)	(8.5)	(2.0)	(0.5)	(0.1)	(0.1)			(100.0)				
合計	2	644	15,344	47,476	56,055	33,948	13,033	4,442	1,612	540	215	138	173,449		173,449	55.20	12.75
(%)	(0.0)	(0.4)	(8.8)	(27.4)	(32.3)	(19.6)	(7.5)	(2.6)	(0.9)	(0.3)	(0.1)	(0.1)	(100.0)				
記載なし	60	12	28	57	51	18	8	9	3				246	4,514	4,760	40.58	21.43
(%)	(24.4)	(4.9)	(11.4)	(23.2)	(20.7)	(7.3)	(3.3)	(3.7)	(1.2)				(100.0)				
総計	62	656	15,372	47,533	56,106	33,966	13,041	4,451	1,615	540	215	138	173,695	4,514	178,209	55.17	12.78
(%)	(0.0)	(0.4)	(8.8)	(27.4)	(32.3)	(19.6)	(7.5)	(2.6)	(0.9)	(0.3)	(0.1)	(0.1)	(100.0)				
平均	7.54	5.04	4.93	4.69	4.49	4.34	4.29	4.28	4.32	4.30	4.24	3.93	4.53		4.53		
標準偏差	3.77	2.48	1.98	1.76	1.60	1.47	1.39	1.31	1.21	1.28	1.19	1.14	1.65		1.65		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査