

表 2491 透析後体重(kg) 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回）

透析前ヘモグロビン濃度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
5未満	1	1	26	82	81	45	15	3	2	1			257	4	261	53.16	12.11
(%)	(0.4)	(0.4)	(10.1)	(31.9)	(31.5)	(17.5)	(5.8)	(1.2)	(0.8)	(0.4)			(100.0)				
5～		2	36	111	124	54	17	5	3	1			353	9	362	52.94	11.89
(%)		(0.6)	(10.2)	(31.4)	(35.1)	(15.3)	(4.8)	(1.4)	(0.8)	(0.3)			(100.0)				
6～		5	112	284	276	158	58	20	4	4		1	922	20	942	53.17	12.66
(%)		(0.5)	(12.1)	(30.8)	(29.9)	(17.1)	(6.3)	(2.2)	(0.4)	(0.4)		(0.1)	(100.0)				
7～	1	36	460	1,136	1,095	514	190	61	22	8	5	4	3,532	46	3,578	52.38	13.14
(%)	(0.0)	(1.0)	(13.0)	(32.2)	(31.0)	(14.6)	(5.4)	(1.7)	(0.6)	(0.2)	(0.1)	(0.1)	(100.0)				
8～	6	66	1,447	3,678	3,571	1,815	688	200	82	27	16	9	11,605	121	11,726	52.86	12.56
(%)	(0.1)	(0.6)	(12.5)	(31.7)	(30.8)	(15.6)	(5.9)	(1.7)	(0.7)	(0.2)	(0.1)	(0.1)	(100.0)				
9～	12	152	3,808	10,944	11,897	6,551	2,191	789	273	93	43	33	36,786	252	37,038	53.91	12.49
(%)	(0.0)	(0.4)	(10.4)	(29.8)	(32.3)	(17.8)	(6.0)	(2.1)	(0.7)	(0.3)	(0.1)	(0.1)	(100.0)				
10～	32	249	6,102	20,242	24,639	14,920	5,646	1,943	683	247	95	57	74,855	362	75,217	55.42	12.65
(%)	(0.0)	(0.3)	(8.2)	(27.0)	(32.9)	(19.9)	(7.5)	(2.6)	(0.9)	(0.3)	(0.1)	(0.1)	(100.0)				
11～	23	174	4,547	15,307	20,050	12,912	5,216	1,868	683	240	88	46	61,154	325	61,479	56.35	12.97
(%)	(0.0)	(0.3)	(7.4)	(25.0)	(32.8)	(21.1)	(8.5)	(3.1)	(1.1)	(0.4)	(0.1)	(0.1)	(100.0)				
12～	1	71	1,673	5,368	7,112	4,766	2,076	765	269	82	31	22	22,236	134	22,370	56.76	13.21
(%)	(0.0)	(0.3)	(7.5)	(24.1)	(32.0)	(21.4)	(9.3)	(3.4)	(1.2)	(0.4)	(0.1)	(0.1)	(100.0)				
13～	5	18	409	1,264	1,637	1,204	590	231	87	33	6	10	5,494	43	5,537	57.75	14.14
(%)	(0.1)	(0.3)	(7.4)	(23.0)	(29.8)	(21.9)	(10.7)	(4.2)	(1.6)	(0.6)	(0.1)	(0.2)	(100.0)				
14～	1	6	133	362	499	474	249	104	59	13	12	5	1,917	16	1,933	60.26	15.40
(%)	(0.1)	(0.3)	(6.9)	(18.9)	(26.0)	(24.7)	(13.0)	(5.4)	(3.1)	(0.7)	(0.6)	(0.3)	(100.0)				
合計	82	780	18,753	58,778	70,981	43,413	16,936	5,989	2,167	749	296	187	219,111	1,332	220,443	55.46	12.90
(%)	(0.0)	(0.4)	(8.6)	(26.8)	(32.4)	(19.8)	(7.7)	(2.7)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
記載なし	2	22	364	1,031	1,128	669	248	92	28	6	3	2	3,595	4,933	8,528	54.30	12.57
(%)	(0.1)	(0.6)	(10.1)	(28.7)	(31.4)	(18.6)	(6.9)	(2.6)	(0.8)	(0.2)	(0.1)	(0.1)	(100.0)				
総計	84	802	19,117	59,809	72,109	44,082	17,184	6,081	2,195	755	299	189	222,706	6,265	228,971	55.45	12.90
(%)	(0.0)	(0.4)	(8.6)	(26.9)	(32.4)	(19.8)	(7.7)	(2.7)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
平均	10.53	10.43	10.49	10.58	10.69	10.80	10.90	10.97	10.99	10.93	10.90	10.85	10.69	10.45	10.69		
標準偏差	1.47	1.47	1.33	1.28	1.25	1.25	1.28	1.28	1.33	1.31	1.40	1.45	1.28	1.57	1.28		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査