

表 2495 透析後体重(kg) 透析前収縮期血圧(mmHg)別（施設血液透析患者 透析回数週3回）

透析前収縮期血圧	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
60未満		2	4	15	10	1		1					33	2	35	47.18	10.98
(%)		(6.1)	(12.1)	(45.5)	(30.3)	(3.0)		(3.0)					(100.0)				
60～		7	59	80	72	31	12	4			1		266	5	271	49.46	12.92
(%)		(2.6)	(22.2)	(30.1)	(27.1)	(11.7)	(4.5)	(1.5)			(0.4)		(100.0)				
80～	3	30	371	799	658	350	103	36	7	4	2	1	2,364	13	2,377	51.05	12.15
(%)	(0.1)	(1.3)	(15.7)	(33.8)	(27.8)	(14.8)	(4.4)	(1.5)	(0.3)	(0.2)	(0.1)	(0.0)	(100.0)				
100～	4	80	1,571	4,038	4,124	2,188	720	247	78	25	11	8	13,094	61	13,155	53.03	12.21
(%)	(0.0)	(0.6)	(12.0)	(30.8)	(31.5)	(16.7)	(5.5)	(1.9)	(0.6)	(0.2)	(0.1)	(0.1)	(100.0)				
120～	12	159	3,956	11,740	13,758	7,650	2,769	887	315	110	34	26	41,416	160	41,576	54.41	12.35
(%)	(0.0)	(0.4)	(9.6)	(28.3)	(33.2)	(18.5)	(6.7)	(2.1)	(0.8)	(0.3)	(0.1)	(0.1)	(100.0)				
140～	26	191	5,083	17,238	20,880	13,157	4,899	1,697	544	173	71	51	64,010	193	64,203	55.52	12.54
(%)	(0.0)	(0.3)	(7.9)	(26.9)	(32.6)	(20.6)	(7.7)	(2.7)	(0.8)	(0.3)	(0.1)	(0.1)	(100.0)				
160～	20	120	3,518	11,780	15,173	9,737	3,998	1,468	564	198	75	41	46,692	151	46,843	56.39	13.15
(%)	(0.0)	(0.3)	(7.5)	(25.2)	(32.5)	(20.9)	(8.6)	(3.1)	(1.2)	(0.4)	(0.2)	(0.1)	(100.0)				
180～	7	68	1,410	4,248	5,525	3,848	1,723	675	281	99	50	19	17,953	49	18,002	57.19	13.98
(%)	(0.0)	(0.4)	(7.9)	(23.7)	(30.8)	(21.4)	(9.6)	(3.8)	(1.6)	(0.6)	(0.3)	(0.1)	(100.0)				
200～	3	11	341	1,091	1,335	906	462	206	86	34	19	9	4,503	12	4,515	57.89	14.79
(%)	(0.1)	(0.2)	(7.6)	(24.2)	(29.6)	(20.1)	(10.3)	(4.6)	(1.9)	(0.8)	(0.4)	(0.2)	(100.0)				
220～		2	73	228	260	156	85	54	22	5	1	1	887	7	894	57.65	14.97
(%)		(0.2)	(8.2)	(25.7)	(29.3)	(17.6)	(9.6)	(6.1)	(2.5)	(0.6)	(0.1)	(0.1)	(100.0)				
合計	75	670	16,386	51,257	61,795	38,024	14,771	5,275	1,897	648	264	156	191,218	653	191,871	55.48	12.90
(%)	(0.0)	(0.4)	(8.6)	(26.8)	(32.3)	(19.9)	(7.7)	(2.8)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
記載なし	9	132	2,731	8,552	10,314	6,058	2,413	806	298	107	35	33	31,488	5,612	37,100	55.23	12.91
(%)	(0.0)	(0.4)	(8.7)	(27.2)	(32.8)	(19.2)	(7.7)	(2.6)	(0.9)	(0.3)	(0.1)	(0.1)	(100.0)				
総計	84	802	19,117	59,809	72,109	44,082	17,184	6,081	2,195	755	299	189	222,706	6,265	228,971	55.45	12.90
(%)	(0.0)	(0.4)	(8.6)	(26.9)	(32.4)	(19.8)	(7.7)	(2.7)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
平均	152.44	144.71	148.13	149.88	151.46	153.16	155.39	157.23	159.33	159.68	161.50	158.73	151.65	148.19	151.64		
標準偏差	24.62	28.14	25.58	24.20	23.51	23.21	23.64	24.44	24.72	24.64	25.91	25.27	24.01	26.45	24.02		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査