

表 2661 体重減少量(kg) %クレアチニン産生速度(%)別(血液透析濾過患者 男性のみ 透析回数週3回 透析歴2年以上)

%クレアチニン 産生速度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
60未満	12	128	307	373	143	34	5				1,002		1,002	2.05	1.04
(%)	(1.2)	(12.8)	(30.6)	(37.2)	(14.3)	(3.4)	(0.5)				(100.0)				
60～	8	79	225	358	159	39	15	1	2		886		886	2.35	1.11
(%)	(0.9)	(8.9)	(25.4)	(40.4)	(17.9)	(4.4)	(1.7)	(0.1)	(0.2)		(100.0)				
70～	9	80	345	638	361	101	29	5	1	1	1,570	1	1,571	2.55	1.08
(%)	(0.6)	(5.1)	(22.0)	(40.6)	(23.0)	(6.4)	(1.8)	(0.3)	(0.1)	(0.1)	(100.0)				
80～	9	93	480	1,070	625	211	32	6	1	2	2,529		2,529	2.65	1.02
(%)	(0.4)	(3.7)	(19.0)	(42.3)	(24.7)	(8.3)	(1.3)	(0.2)	(0.0)	(0.1)	(100.0)				
90～	2	76	558	1,285	918	305	58	5			3,207	1	3,208	2.78	0.98
(%)	(0.1)	(2.4)	(17.4)	(40.1)	(28.6)	(9.5)	(1.8)	(0.2)			(100.0)				
100～	6	61	538	1,415	1,122	358	75	11	3	3	3,592		3,592	2.88	1.03
(%)	(0.2)	(1.7)	(15.0)	(39.4)	(31.2)	(10.0)	(2.1)	(0.3)	(0.1)	(0.1)	(100.0)				
110～	4	44	397	1,282	1,189	363	57	9	3	4	3,352		3,352	2.94	0.97
(%)	(0.1)	(1.3)	(11.8)	(38.2)	(35.5)	(10.8)	(1.7)	(0.3)	(0.1)	(0.1)	(100.0)				
120～	3	35	250	940	780	253	54	5		1	2,321		2,321	2.97	1.01
(%)	(0.1)	(1.5)	(10.8)	(40.5)	(33.6)	(10.9)	(2.3)	(0.2)		(0.0)	(100.0)				
130～	1	17	131	486	432	169	16	6	1	4	1,263		1,263	3.01	1.07
(%)	(0.1)	(1.3)	(10.4)	(38.5)	(34.2)	(13.4)	(1.3)	(0.5)	(0.1)	(0.3)	(100.0)				
140～		8	74	319	311	97	20	1	2	6	838	2	840	3.14	1.57
(%)		(1.0)	(8.8)	(38.1)	(37.1)	(11.6)	(2.4)	(0.1)	(0.2)	(0.7)	(100.0)				
合計	54	621	3,305	8,166	6,040	1,930	361	49	13	21	20,560	4	20,564	2.78	1.07
(%)	(0.3)	(3.0)	(16.1)	(39.7)	(29.4)	(9.4)	(1.8)	(0.2)	(0.1)	(0.1)	(100.0)				
記載なし	4	40	136	284	183	48	14	3		1	713	499	1,212	2.61	1.10
(%)	(0.6)	(5.6)	(19.1)	(39.8)	(25.7)	(6.7)	(2.0)	(0.4)		(0.1)	(100.0)				
総計	58	661	3,441	8,450	6,223	1,978	375	52	13	22	21,273	503	21,776	2.78	1.08
(%)	(0.3)	(3.1)	(16.2)	(39.7)	(29.3)	(9.3)	(1.8)	(0.2)	(0.1)	(0.1)	(100.0)				
平均	78.04	81.98	93.70	101.55	106.35	107.19	105.30	104.93	107.84	126.82	101.68	156.48	101.69		
標準偏差	27.40	28.04	24.97	23.26	21.92	21.06	21.38	19.43	32.25	28.38	23.82	103.64	23.86		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査