

表 2709 体重減少率(%) HDL-C濃度(mg/dL)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

HDL-C濃度	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
20未満	37	57	98	99	116	64	22	6	4		503	11	514	3.53	1.71
(%)	(7.4)	(11.3)	(19.5)	(19.7)	(23.1)	(12.7)	(4.4)	(1.2)	(0.8)		(100.0)				
20～	260	471	1,215	2,420	2,887	1,678	639	204	45	31	9,850	83	9,933	4.16	1.52
(%)	(2.6)	(4.8)	(12.3)	(24.6)	(29.3)	(17.0)	(6.5)	(2.1)	(0.5)	(0.3)	(100.0)				
30～	590	1,140	3,003	6,429	8,155	5,571	2,330	733	191	118	28,260	218	28,478	4.34	1.52
(%)	(2.1)	(4.0)	(10.6)	(22.7)	(28.9)	(19.7)	(8.2)	(2.6)	(0.7)	(0.4)	(100.0)				
40～	525	1,188	3,036	6,326	8,802	6,676	3,296	1,196	328	177	31,550	228	31,778	4.53	1.58
(%)	(1.7)	(3.8)	(9.6)	(20.1)	(27.9)	(21.2)	(10.4)	(3.8)	(1.0)	(0.6)	(100.0)				
50～	420	719	2,015	4,241	5,976	5,147	2,661	1,065	338	184	22,766	139	22,905	4.67	1.65
(%)	(1.8)	(3.2)	(8.9)	(18.6)	(26.2)	(22.6)	(11.7)	(4.7)	(1.5)	(0.8)	(100.0)				
60～	247	399	1,015	2,195	3,373	3,023	1,785	760	257	145	13,199	79	13,278	4.82	1.71
(%)	(1.9)	(3.0)	(7.7)	(16.6)	(25.6)	(22.9)	(13.5)	(5.8)	(1.9)	(1.1)	(100.0)				
70～	132	196	475	1,062	1,530	1,509	977	427	139	82	6,529	46	6,575	4.90	1.76
(%)	(2.0)	(3.0)	(7.3)	(16.3)	(23.4)	(23.1)	(15.0)	(6.5)	(2.1)	(1.3)	(100.0)				
80～	58	88	222	483	749	710	467	212	86	37	3,112	13	3,125	4.95	1.78
(%)	(1.9)	(2.8)	(7.1)	(15.5)	(24.1)	(22.8)	(15.0)	(6.8)	(2.8)	(1.2)	(100.0)				
90～	41	47	100	207	332	314	186	106	41	22	1,396	12	1,408	4.91	1.87
(%)	(2.9)	(3.4)	(7.2)	(14.8)	(23.8)	(22.5)	(13.3)	(7.6)	(2.9)	(1.6)	(100.0)				
100～	33	38	87	179	271	269	152	79	30	14	1,152	9	1,161	4.84	1.87
(%)	(2.9)	(3.3)	(7.6)	(15.5)	(23.5)	(23.4)	(13.2)	(6.9)	(2.6)	(1.2)	(100.0)				
合計	2,343	4,343	11,266	23,641	32,191	24,961	12,515	4,788	1,459	810	118,317	838	119,155	4.55	1.63
(%)	(2.0)	(3.7)	(9.5)	(20.0)	(27.2)	(21.1)	(10.6)	(4.0)	(1.2)	(0.7)	(100.0)				
記載なし	1,353	2,266	5,655	10,943	14,224	11,328	5,912	2,320	736	395	55,132	3,922	59,054	4.51	1.69
(%)	(2.5)	(4.1)	(10.3)	(19.8)	(25.8)	(20.5)	(10.7)	(4.2)	(1.3)	(0.7)	(100.0)				
総計	3,696	6,609	16,921	34,584	46,415	36,289	18,427	7,108	2,195	1,205	173,449	4,760	178,209	4.53	1.65
(%)	(2.1)	(3.8)	(9.8)	(19.9)	(26.8)	(20.9)	(10.6)	(4.1)	(1.3)	(0.7)	(100.0)				
平均	47.76	46.15	46.06	46.41	47.51	49.67	51.96	54.18	56.17	55.50	48.47	46.91	48.46		
標準偏差	18.25	16.64	16.06	15.84	16.20	16.57	16.99	17.55	18.02	17.63	16.61	17.78	16.62		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査