

表 2785 Body Mass Index 透析前クレアチニン濃度(mg/dL)別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前クレアチニン濃度	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
2未満	4	11	11	13	8	7	2		3	2	61	15	76	19.21	4.42
(%)	(6.6)	(18.0)	(18.0)	(21.3)	(13.1)	(11.5)	(3.3)		(4.9)	(3.3)	(100.0)				
2～	117	221	354	290	213	108	61	31	18	9	1,422	234	1,656	18.59	3.65
(%)	(8.2)	(15.5)	(24.9)	(20.4)	(15.0)	(7.6)	(4.3)	(2.2)	(1.3)	(0.6)	(100.0)				
4～	357	1,125	1,686	1,566	1,006	604	287	154	75	50	6,910	885	7,795	18.80	3.54
(%)	(5.2)	(16.3)	(24.4)	(22.7)	(14.6)	(8.7)	(4.2)	(2.2)	(1.1)	(0.7)	(100.0)				
6～	312	1,995	4,532	5,532	4,421	2,677	1,338	679	324	226	22,036	1,970	24,006	19.92	3.43
(%)	(1.4)	(9.1)	(20.6)	(25.1)	(20.1)	(12.1)	(6.1)	(3.1)	(1.5)	(1.0)	(100.0)				
8～	195	1,835	6,607	10,803	10,062	6,833	3,438	1,827	853	703	43,156	2,959	46,115	20.82	3.41
(%)	(0.5)	(4.3)	(15.3)	(25.0)	(23.3)	(15.8)	(8.0)	(4.2)	(2.0)	(1.6)	(100.0)				
10～	76	870	4,692	9,952	11,538	8,465	4,969	2,490	1,218	984	45,254	2,780	48,034	21.61	3.39
(%)	(0.2)	(1.9)	(10.4)	(22.0)	(25.5)	(18.7)	(11.0)	(5.5)	(2.7)	(2.2)	(100.0)				
12～	15	211	1,619	4,830	6,925	6,284	4,088	2,087	1,069	973	28,101	1,653	29,754	22.53	3.44
(%)	(0.1)	(0.8)	(5.8)	(17.2)	(24.6)	(22.4)	(14.5)	(7.4)	(3.8)	(3.5)	(100.0)				
14～	9	30	372	1,365	2,536	2,639	1,948	1,174	682	575	11,330	706	12,036	23.43	3.54
(%)	(0.1)	(0.3)	(3.3)	(12.0)	(22.4)	(23.3)	(17.2)	(10.4)	(6.0)	(5.1)	(100.0)				
16～	1	4	71	309	724	794	732	521	319	325	3,800	300	4,100	24.40	3.70
(%)	(0.0)	(0.1)	(1.9)	(8.1)	(19.1)	(20.9)	(19.3)	(13.7)	(8.4)	(8.6)	(100.0)				
合計	1,086	6,302	19,944	34,660	37,433	28,411	16,863	8,963	4,561	3,847	162,070	11,502	173,572	21.38	3.65
(%)	(0.7)	(3.9)	(12.3)	(21.4)	(23.1)	(17.5)	(10.4)	(5.5)	(2.8)	(2.4)	(100.0)				
記載なし	10	27	105	156	137	131	57	42	13	17	695	3,942	4,637	21.11	3.70
(%)	(1.4)	(3.9)	(15.1)	(22.4)	(19.7)	(18.8)	(8.2)	(6.0)	(1.9)	(2.4)	(100.0)				
総計	1,096	6,329	20,049	34,816	37,570	28,542	16,920	9,005	4,574	3,864	162,765	15,444	178,209	21.37	3.65
(%)	(0.7)	(3.9)	(12.3)	(21.4)	(23.1)	(17.5)	(10.4)	(5.5)	(2.8)	(2.4)	(100.0)				
平均	6.68	7.82	8.99	9.84	10.50	10.93	11.32	11.48	11.71	11.91	10.33	9.83	10.29		
標準偏差	2.61	2.28	2.47	2.47	2.58	2.61	2.70	2.83	2.94	2.88	2.76	3.05	2.79		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査