

表 2836 Body Mass Index 透析前リン濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

透析前リン濃度	14未満	14～	16～	18～	20～	22～	24～	26～	28～	30～	合計	記載なし	総計	平均	標準偏差
3未満	21	71	177	248	216	140	89	56	15	14	1,047	86	1,133	20.47	3.60
(%)	(2.0)	(6.8)	(16.9)	(23.7)	(20.6)	(13.4)	(8.5)	(5.3)	(1.4)	(1.3)	(100.0)				
3.0～	5	48	213	325	371	255	138	50	31	23	1,459	115	1,574	21.00	3.35
(%)	(0.3)	(3.3)	(14.6)	(22.3)	(25.4)	(17.5)	(9.5)	(3.4)	(2.1)	(1.6)	(100.0)				
3.5～	9	99	365	581	679	414	260	148	49	44	2,648	201	2,849	21.13	3.44
(%)	(0.3)	(3.7)	(13.8)	(21.9)	(25.6)	(15.6)	(9.8)	(5.6)	(1.9)	(1.7)	(100.0)				
4.0～	14	139	514	937	1,018	776	444	213	110	90	4,255	310	4,565	21.38	3.48
(%)	(0.3)	(3.3)	(12.1)	(22.0)	(23.9)	(18.2)	(10.4)	(5.0)	(2.6)	(2.1)	(100.0)				
4.5～	26	142	555	1,187	1,313	1,004	585	279	147	121	5,359	353	5,712	21.51	3.47
(%)	(0.5)	(2.6)	(10.4)	(22.1)	(24.5)	(18.7)	(10.9)	(5.2)	(2.7)	(2.3)	(100.0)				
5.0～	10	167	651	1,188	1,343	1,072	631	337	192	144	5,735	399	6,134	21.65	3.58
(%)	(0.2)	(2.9)	(11.4)	(20.7)	(23.4)	(18.7)	(11.0)	(5.9)	(3.3)	(2.5)	(100.0)				
5.5～	19	116	542	1,030	1,080	917	585	329	168	145	4,931	368	5,299	21.81	3.67
(%)	(0.4)	(2.4)	(11.0)	(20.9)	(21.9)	(18.6)	(11.9)	(6.7)	(3.4)	(2.9)	(100.0)				
6.0～	1	31	95	206	191	159	95	61	21	21	881	69	950	21.53	3.56
(%)	(0.1)	(3.5)	(10.8)	(23.4)	(21.7)	(18.0)	(10.8)	(6.9)	(2.4)	(2.4)	(100.0)				
6.1～	13	76	308	627	670	531	335	207	113	97	2,977	267	3,244	21.86	3.77
(%)	(0.4)	(2.6)	(10.3)	(21.1)	(22.5)	(17.8)	(11.3)	(7.0)	(3.8)	(3.3)	(100.0)				
6.5～	12	49	231	503	602	470	342	178	120	83	2,590	216	2,806	22.12	3.71
(%)	(0.5)	(1.9)	(8.9)	(19.4)	(23.2)	(18.1)	(13.2)	(6.9)	(4.6)	(3.2)	(100.0)				
7.0～	6	48	116	323	360	283	228	128	63	85	1,640	135	1,775	22.35	3.93
(%)	(0.4)	(2.9)	(7.1)	(19.7)	(22.0)	(17.3)	(13.9)	(7.8)	(3.8)	(5.2)	(100.0)				
7.5～	5	24	71	160	240	223	129	76	50	49	1,027	77	1,104	22.51	3.83
(%)	(0.5)	(2.3)	(6.9)	(15.6)	(23.4)	(21.7)	(12.6)	(7.4)	(4.9)	(4.8)	(100.0)				
8.0～	2	10	62	109	122	105	71	44	27	36	588	63	651	22.36	4.05
(%)	(0.3)	(1.7)	(10.5)	(18.5)	(20.7)	(17.9)	(12.1)	(7.5)	(4.6)	(6.1)	(100.0)				
8.5～	2	4	34	67	68	78	61	32	21	27	394	29	423	22.92	4.11
(%)	(0.5)	(1.0)	(8.6)	(17.0)	(17.3)	(19.8)	(15.5)	(8.1)	(5.3)	(6.9)	(100.0)				
9.0～	3	11	47	85	138	108	82	54	44	37	609	73	682	22.95	4.17
(%)	(0.5)	(1.8)	(7.7)	(14.0)	(22.7)	(17.7)	(13.5)	(8.9)	(7.2)	(6.1)	(100.0)				
合計	148	1,035	3,981	7,576	8,411	6,535	4,075	2,192	1,171	1,016	36,140	2,761	38,901	21.67	3.65
(%)	(0.4)	(2.9)	(11.0)	(21.0)	(23.3)	(18.1)	(11.3)	(6.1)	(3.2)	(2.8)	(100.0)				
記載なし	1	11	17	38	51	36	13	5	6	1	179	866	1,045	20.99	3.24
(%)	(0.6)	(6.1)	(9.5)	(21.2)	(28.5)	(20.1)	(7.3)	(2.8)	(3.4)	(0.6)	(100.0)				
総計	149	1,046	3,998	7,614	8,462	6,571	4,088	2,197	1,177	1,017	36,319	3,627	39,946	21.67	3.65
(%)	(0.4)	(2.9)	(11.0)	(21.0)	(23.3)	(18.1)	(11.3)	(6.0)	(3.2)	(2.8)	(100.0)				
平均	5.04	5.03	5.12	5.25	5.30	5.40	5.49	5.57	5.76	5.88	5.35	5.45	5.36		
標準偏差	1.75	1.45	1.39	1.35	1.41	1.39	1.43	1.45	1.51	1.55	1.42	1.54	1.43		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査