

表 3223 透析前アルブミン濃度(g/dL) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
施設血液透析	159	852	3,435	12,892	54,438	111,197	42,314	2,504	178	227,969	27,672	255,641	3.60	0.45
(%)	(0.1)	(0.4)	(1.5)	(5.7)	(23.9)	(48.8)	(18.6)	(1.1)	(0.1)	(100.0)				
Off-line HDF	5	24	96	341	1,471	2,899	917	35	4	5,792	523	6,315	3.57	0.44
(%)	(0.1)	(0.4)	(1.7)	(5.9)	(25.4)	(50.1)	(15.8)	(0.6)	(0.1)	(100.0)				
On-line HDF	6	40	206	1,050	7,335	17,840	6,334	296	15	33,122	2,968	36,090	3.65	0.38
(%)	(0.0)	(0.1)	(0.6)	(3.2)	(22.1)	(53.9)	(19.1)	(0.9)	(0.0)	(100.0)				
Push/Pull HDF		1	1	23	130	236	75			466	71	537	3.59	0.38
(%)		(0.2)	(0.2)	(4.9)	(27.9)	(50.6)	(16.1)			(100.0)				
ハイオフィльтраション		1	6	22	90	156	47	1		323	18	341	3.53	0.43
(%)		(0.3)	(1.9)	(6.8)	(27.9)	(48.3)	(14.6)	(0.3)		(100.0)				
血液濾過			3	2	21	48	12	2	1	89	3	92	3.61	0.46
(%)			(3.4)	(2.2)	(23.6)	(53.9)	(13.5)	(2.2)	(1.1)	(100.0)				
血液吸着透析 (リクセル)			15	63	425	808	220	7	2	1,540	83	1,623	3.59	0.38
(%)			(1.0)	(4.1)	(27.6)	(52.5)	(14.3)	(0.5)	(0.1)	(100.0)				
血液吸着透析 (その他)					1	2	1			4	3	7	3.75	0.25
(%)					(25.0)	(50.0)	(25.0)			(100.0)				
在宅血液透析			1	6	39	143	157	48	1	395	126	521	3.95	0.42
(%)			(0.3)	(1.5)	(9.9)	(36.2)	(39.7)	(12.2)	(0.3)	(100.0)				
腹膜透析 (CAPD)	10	40	166	563	1,149	1,056	220	6	1	3,211	1,810	5,021	3.27	0.52
(%)	(0.3)	(1.2)	(5.2)	(17.5)	(35.8)	(32.9)	(6.9)	(0.2)	(0.0)	(100.0)				
腹膜透析 (APD)	5	27	108	364	687	594	157	11		1,953	886	2,839	3.27	0.54
(%)	(0.3)	(1.4)	(5.5)	(18.6)	(35.2)	(30.4)	(8.0)	(0.6)		(100.0)				
腹膜透析 (CCPD)		4	24	110	322	256	54	4	1	775	306	1,081	3.34	0.47
(%)		(0.5)	(3.1)	(14.2)	(41.5)	(33.0)	(7.0)	(0.5)	(0.1)	(100.0)				
合計	185	989	4,061	15,436	66,108	135,235	50,508	2,914	203	275,639	34,469	310,108	3.60	0.44
(%)	(0.1)	(0.4)	(1.5)	(5.6)	(24.0)	(49.1)	(18.3)	(1.1)	(0.1)	(100.0)				
記載なし														
(%)														
総計	185	989	4,061	15,436	66,108	135,235	50,508	2,914	203	275,639	34,469	310,108	3.60	0.44
(%)	(0.1)	(0.4)	(1.5)	(5.6)	(24.0)	(49.1)	(18.3)	(1.1)	(0.1)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2014年調査