

表 3362 CRP濃度(mg/dL) 総コレステロール濃度(mg/dL)別（腹膜透析患者）

総コレステロール濃度	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
100未満	23	8	3	3	4	1	2	1	1	2	48	2	50	2.51	6.52
(%)	(47.9)	(16.7)	(6.3)	(6.3)	(8.3)	(2.1)	(4.2)	(2.1)	(2.1)	(4.2)	(100.0)				
100～	68	28	16	12	9	4	2		2		141	19	160	0.92	1.99
(%)	(48.2)	(19.9)	(11.3)	(8.5)	(6.4)	(2.8)	(1.4)		(1.4)		(100.0)				
120～	228	80	36	17	16	10	1	3	3	2	396	52	448	0.75	2.26
(%)	(57.6)	(20.2)	(9.1)	(4.3)	(4.0)	(2.5)	(0.3)	(0.8)	(0.8)	(0.5)	(100.0)				
140～	396	154	81	37	26	15	9	3	3		724	82	806	0.62	1.43
(%)	(54.7)	(21.3)	(11.2)	(5.1)	(3.6)	(2.1)	(1.2)	(0.4)	(0.4)		(100.0)				
160～	458	152	61	48	31	11	4	4	6		775	90	865	0.60	1.51
(%)	(59.1)	(19.6)	(7.9)	(6.2)	(4.0)	(1.4)	(0.5)	(0.5)	(0.8)		(100.0)				
180～	480	143	58	38	23	5	9	2	2		760	73	833	0.47	1.22
(%)	(63.2)	(18.8)	(7.6)	(5.0)	(3.0)	(0.7)	(1.2)	(0.3)	(0.3)		(100.0)				
200～	311	96	35	31	18	5		2	4		502	62	564	0.50	1.30
(%)	(62.0)	(19.1)	(7.0)	(6.2)	(3.6)	(1.0)		(0.4)	(0.8)		(100.0)				
220～	198	41	19	15	7	2			1		283	35	318	0.36	0.92
(%)	(70.0)	(14.5)	(6.7)	(5.3)	(2.5)	(0.7)			(0.4)		(100.0)				
240～	92	24	12	8	5	2	1				144	15	159	0.44	0.98
(%)	(63.9)	(16.7)	(8.3)	(5.6)	(3.5)	(1.4)	(0.7)				(100.0)				
260～	80	25	13	3	7	2		2			132	17	149	0.55	1.37
(%)	(60.6)	(18.9)	(9.8)	(2.3)	(5.3)	(1.5)		(1.5)			(100.0)				
合計	2,334	751	334	212	146	57	28	17	22	4	3,905	447	4,352	0.59	1.65
(%)	(59.8)	(19.2)	(8.6)	(5.4)	(3.7)	(1.5)	(0.7)	(0.4)	(0.6)	(0.1)	(100.0)				
記載なし	733	244	115	96	56	26	12	7	12	3	1,304	3,285	4,589	0.74	1.88
(%)	(56.2)	(18.7)	(8.8)	(7.4)	(4.3)	(2.0)	(0.9)	(0.5)	(0.9)	(0.2)	(100.0)				
総計	3,067	995	449	308	202	83	40	24	34	7	5,209	3,732	8,941	0.63	1.71
(%)	(58.9)	(19.1)	(8.6)	(5.9)	(3.9)	(1.6)	(0.8)	(0.5)	(0.7)	(0.1)	(100.0)				
平均	181.48	176.99	175.81	176.26	175.58	165.61	159.39	178.76	165.32	103.75	179.05	178.66	179.01		
標準偏差	40.31	39.45	41.00	37.89	44.48	40.65	35.28	60.23	37.39	31.70	40.48	41.09	40.54		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査