

表 3386 透析前補正カルシウム濃度(mg/dL) 透析前脈拍(拍/分)別 (透析患者全体)

透析前脈拍	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
40未満	1		6	16	4	44	47	34	6	14	3	3	2	1	181	4	185	9.20	0.87
(%)	(0.6)		(3.3)	(8.8)	(2.2)	(24.3)	(26.0)	(18.8)	(3.3)	(7.7)	(1.7)	(1.7)	(1.1)	(0.6)	(100.0)				
40～	6	12	52	168	45	580	705	469	42	148	62	18	16	3	2,326	55	2,381	9.19	0.72
(%)	(0.3)	(0.5)	(2.2)	(7.2)	(1.9)	(24.9)	(30.3)	(20.2)	(1.8)	(6.4)	(2.7)	(0.8)	(0.7)	(0.1)	(100.0)				
50～	62	87	399	1,497	414	4,920	5,800	3,643	386	1,077	537	179	64	40	19,105	407	19,512	9.17	0.72
(%)	(0.3)	(0.5)	(2.1)	(7.8)	(2.2)	(25.8)	(30.4)	(19.1)	(2.0)	(5.6)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
60～	210	318	1,349	4,540	1,317	15,059	17,977	11,419	1,420	3,466	1,642	572	187	126	59,602	1,179	60,781	9.17	0.73
(%)	(0.4)	(0.5)	(2.3)	(7.6)	(2.2)	(25.3)	(30.2)	(19.2)	(2.4)	(5.8)	(2.8)	(1.0)	(0.3)	(0.2)	(100.0)				
70～	330	474	1,618	5,247	1,542	17,903	21,528	14,013	1,722	4,385	2,065	644	225	156	71,852	1,420	73,272	9.18	0.74
(%)	(0.5)	(0.7)	(2.3)	(7.3)	(2.1)	(24.9)	(30.0)	(19.5)	(2.4)	(6.1)	(2.9)	(0.9)	(0.3)	(0.2)	(100.0)				
80～	197	269	1,052	3,387	944	10,911	13,748	9,174	1,152	2,869	1,399	456	162	111	45,831	983	46,814	9.20	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(23.8)	(30.0)	(20.0)	(2.5)	(6.3)	(3.1)	(1.0)	(0.4)	(0.2)	(100.0)				
90～	74	130	462	1,404	408	4,747	5,959	4,102	512	1,369	672	222	86	58	20,205	385	20,590	9.22	0.76
(%)	(0.4)	(0.6)	(2.3)	(6.9)	(2.0)	(23.5)	(29.5)	(20.3)	(2.5)	(6.8)	(3.3)	(1.1)	(0.4)	(0.3)	(100.0)				
100～	28	30	154	389	137	1,331	1,827	1,380	172	396	256	90	32	23	6,245	148	6,393	9.27	0.77
(%)	(0.4)	(0.5)	(2.5)	(6.2)	(2.2)	(21.3)	(29.3)	(22.1)	(2.8)	(6.3)	(4.1)	(1.4)	(0.5)	(0.4)	(100.0)				
110～	4	11	26	90	27	323	434	317	37	129	56	26	10	6	1,496	32	1,528	9.31	0.79
(%)	(0.3)	(0.7)	(1.7)	(6.0)	(1.8)	(21.6)	(29.0)	(21.2)	(2.5)	(8.6)	(3.7)	(1.7)	(0.7)	(0.4)	(100.0)				
120～	3	9	17	45	6	127	172	130	15	67	31	11	2	4	639	16	655	9.30	0.86
(%)	(0.5)	(1.4)	(2.7)	(7.0)	(0.9)	(19.9)	(26.9)	(20.3)	(2.3)	(10.5)	(4.9)	(1.7)	(0.3)	(0.6)	(100.0)				
合計	915	1,340	5,135	16,783	4,844	55,945	68,197	44,681	5,464	13,920	6,723	2,221	786	528	227,482	4,629	232,111	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.6)	(30.0)	(19.6)	(2.4)	(6.1)	(3.0)	(1.0)	(0.3)	(0.2)	(100.0)				
記載なし	227	309	1,264	3,656	1,028	11,374	13,807	9,288	1,134	3,057	1,547	560	187	153	47,591	30,406	77,997	9.19	0.77
(%)	(0.5)	(0.6)	(2.7)	(7.7)	(2.2)	(23.9)	(29.0)	(19.5)	(2.4)	(6.4)	(3.3)	(1.2)	(0.4)	(0.3)	(100.0)				
総計	1,142	1,649	6,399	20,439	5,872	67,319	82,004	53,969	6,598	16,977	8,270	2,781	973	681	275,073	35,035	310,108	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
平均	75.49	75.85	74.84	74.23	74.27	74.21	74.58	75.03	75.34	75.38	75.80	75.72	75.82	76.52	74.69	74.79	74.69		
標準偏差	12.53	12.54	12.79	12.57	12.54	12.59	12.78	12.96	12.78	13.20	13.61	13.83	15.24	13.95	12.83	13.08	12.84		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査