

表 3387 透析前補正カルシウム濃度(mg/dL) 性別（施設血液透析患者 透析回数週3回）

性別	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
男性	549	878	3,537	11,743	3,347	36,800	41,531	25,521	3,055	7,634	3,621	1,152	407	288	140,063	6,970	147,033	9.13	0.73
(%)	(0.4)	(0.6)	(2.5)	(8.4)	(2.4)	(26.3)	(29.7)	(18.2)	(2.2)	(5.5)	(2.6)	(0.8)	(0.3)	(0.2)	(100.0)				
女性	325	388	1,434	4,547	1,329	17,102	23,705	16,883	2,172	5,643	2,832	1,013	362	251	77,986	3,952	81,938	9.28	0.75
(%)	(0.4)	(0.5)	(1.8)	(5.8)	(1.7)	(21.9)	(30.4)	(21.6)	(2.8)	(7.2)	(3.6)	(1.3)	(0.5)	(0.3)	(100.0)				
合計	874	1,266	4,971	16,290	4,676	53,902	65,236	42,404	5,227	13,277	6,453	2,165	769	539	218,049	10,922	228,971	9.18	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.1)	(24.7)	(29.9)	(19.4)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
記載なし																			
(%)																			
総計	874	1,266	4,971	16,290	4,676	53,902	65,236	42,404	5,227	13,277	6,453	2,165	769	539	218,049	10,922	228,971	9.18	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.1)	(24.7)	(29.9)	(19.4)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査