

表 3611 透析前ヘモグロビン濃度(g/dL) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
施設血液透析	269	380	1,011	3,822	12,410	38,710	78,261	64,036	23,322	5,790	2,014	230,025	25,616	255,641	10.68	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.7)	(5.4)	(16.8)	(34.0)	(27.8)	(10.1)	(2.5)	(0.9)	(100.0)				
Off-line HDF	10	10	44	112	356	981	1,858	1,587	627	149	85	5,819	496	6,315	10.68	1.39
(%)	(0.2)	(0.2)	(0.8)	(1.9)	(6.1)	(16.9)	(31.9)	(27.3)	(10.8)	(2.6)	(1.5)	(100.0)				
On-line HDF	16	22	71	343	1,290	4,912	11,163	10,028	3,776	1,045	450	33,116	2,974	36,090	10.87	1.24
(%)	(0.0)	(0.1)	(0.2)	(1.0)	(3.9)	(14.8)	(33.7)	(30.3)	(11.4)	(3.2)	(1.4)	(100.0)				
Push/Pull HDF			2	4	25	86	163	152	60	17	10	519	18	537	10.88	1.29
(%)			(0.4)	(0.8)	(4.8)	(16.6)	(31.4)	(29.3)	(11.6)	(3.3)	(1.9)	(100.0)				
ハイオフィルトレーション		1	1	7	20	34	117	104	34	2	4	324	17	341	10.75	1.23
(%)		(0.3)	(0.3)	(2.2)	(6.2)	(10.5)	(36.1)	(32.1)	(10.5)	(0.6)	(1.2)	(100.0)				
血液濾過				2	5	19	24	27	11	2	1	91	1	92	10.72	1.27
(%)				(2.2)	(5.5)	(20.9)	(26.4)	(29.7)	(12.1)	(2.2)	(1.1)	(100.0)				
血液吸着透析 (リクセル)		2	3	20	93	295	541	412	149	32	7	1,554	69	1,623	10.62	1.19
(%)		(0.1)	(0.2)	(1.3)	(6.0)	(19.0)	(34.8)	(26.5)	(9.6)	(2.1)	(0.5)	(100.0)				
血液吸着透析 (その他)						1	1	1		1		4	3	7	11.27	1.42
(%)						(25.0)	(25.0)	(25.0)		(25.0)		(100.0)				
在宅血液透析	1			1	11	48	78	117	79	33	23	391	130	521	11.45	1.51
(%)	(0.3)			(0.3)	(2.8)	(12.3)	(19.9)	(29.9)	(20.2)	(8.4)	(5.9)	(100.0)				
腹膜透析 (CAPD)	6	11	15	63	188	473	922	925	428	130	50	3,211	1,810	5,021	10.82	1.46
(%)	(0.2)	(0.3)	(0.5)	(2.0)	(5.9)	(14.7)	(28.7)	(28.8)	(13.3)	(4.0)	(1.6)	(100.0)				
腹膜透析 (APD)	1	2	9	45	109	310	560	559	270	81	16	1,962	877	2,839	10.80	1.37
(%)	(0.1)	(0.1)	(0.5)	(2.3)	(5.6)	(15.8)	(28.5)	(28.5)	(13.8)	(4.1)	(0.8)	(100.0)				
腹膜透析 (CCPD)		1	4	16	47	120	224	225	98	26	13	774	307	1,081	10.80	1.36
(%)		(0.1)	(0.5)	(2.1)	(6.1)	(15.5)	(28.9)	(29.1)	(12.7)	(3.4)	(1.7)	(100.0)				
合計	303	429	1,160	4,435	14,554	45,989	93,912	78,173	28,854	7,308	2,673	277,790	32,318	310,108	10.71	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.2)	(16.6)	(33.8)	(28.1)	(10.4)	(2.6)	(1.0)	(100.0)				
記載なし																
(%)																
総計	303	429	1,160	4,435	14,554	45,989	93,912	78,173	28,854	7,308	2,673	277,790	32,318	310,108	10.71	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.2)	(16.6)	(33.8)	(28.1)	(10.4)	(2.6)	(1.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2014年調査