

表 3617 透析前ヘモグロビン濃度(g/dL) 医療機関別 (透析患者全体)

医療機関	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
国公立大学	2	4	6	35	99	154	273	264	91	17	8	953	304	1,257	10.47	1.44
(%)	(0.2)	(0.4)	(0.6)	(3.7)	(10.4)	(16.2)	(28.6)	(27.7)	(9.5)	(1.8)	(0.8)	(100.0)				
私立大学	6	3	17	87	185	436	721	715	271	81	19	2,541	976	3,517	10.58	1.44
(%)	(0.2)	(0.1)	(0.7)	(3.4)	(7.3)	(17.2)	(28.4)	(28.1)	(10.7)	(3.2)	(0.7)	(100.0)				
国立	4	3	12	29	64	127	179	152	53	21	4	648	221	869	10.31	1.59
(%)	(0.6)	(0.5)	(1.9)	(4.5)	(9.9)	(19.6)	(27.6)	(23.5)	(8.2)	(3.2)	(0.6)	(100.0)				
県市町村立	11	27	115	394	1,026	2,770	5,121	4,102	1,657	485	174	15,882	3,303	19,185	10.64	1.37
(%)	(0.1)	(0.2)	(0.7)	(2.5)	(6.5)	(17.4)	(32.2)	(25.8)	(10.4)	(3.1)	(1.1)	(100.0)				
社会保険	3	6	26	87	218	579	1,051	923	363	99	41	3,396	476	3,872	10.66	1.40
(%)	(0.1)	(0.2)	(0.8)	(2.6)	(6.4)	(17.0)	(30.9)	(27.2)	(10.7)	(2.9)	(1.2)	(100.0)				
厚生連	2	8	47	150	447	1,332	2,638	2,201	1,012	243	97	8,177	1,087	9,264	10.76	1.33
(%)	(0.0)	(0.1)	(0.6)	(1.8)	(5.5)	(16.3)	(32.3)	(26.9)	(12.4)	(3.0)	(1.2)	(100.0)				
その他公的	11	46	65	192	581	1,366	2,586	2,567	1,060	310	129	8,913	2,076	10,989	10.73	1.46
(%)	(0.1)	(0.5)	(0.7)	(2.2)	(6.5)	(15.3)	(29.0)	(28.8)	(11.9)	(3.5)	(1.4)	(100.0)				
私立総合	2	25	51	128	325	1,011	2,020	1,871	717	222	62	6,434	1,120	7,554	10.73	1.36
(%)	(0.0)	(0.4)	(0.8)	(2.0)	(5.1)	(15.7)	(31.4)	(29.1)	(11.1)	(3.5)	(1.0)	(100.0)				
私立	80	148	424	1,612	4,936	13,379	25,188	20,975	8,341	2,232	801	78,116	10,526	88,642	10.67	1.35
(%)	(0.1)	(0.2)	(0.5)	(2.1)	(6.3)	(17.1)	(32.2)	(26.9)	(10.7)	(2.9)	(1.0)	(100.0)				
私立診療所	182	159	397	1,721	6,673	24,835	54,135	44,403	15,289	3,598	1,338	152,730	12,229	164,959	10.74	1.21
(%)	(0.1)	(0.1)	(0.3)	(1.1)	(4.4)	(16.3)	(35.4)	(29.1)	(10.0)	(2.4)	(0.9)	(100.0)				
総計	303	429	1,160	4,435	14,554	45,989	93,912	78,173	28,854	7,308	2,673	277,790	32,318	310,108	10.71	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.2)	(16.6)	(33.8)	(28.1)	(10.4)	(2.6)	(1.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査