

表 3619 透析前ヘモグロビン濃度(g/dL) 総コレステロール濃度(mg/dL)別（透析患者全体）

総コレステロール 濃度	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
100未満	14	32	122	377	823	1,623	2,364	1,669	555	145	64	7,788	42	7,830	10.24	1.48
(%)	(0.2)	(0.4)	(1.6)	(4.8)	(10.6)	(20.8)	(30.4)	(21.4)	(7.1)	(1.9)	(0.8)	(100.0)				
100～	17	59	150	616	1,651	4,198	7,676	5,623	1,977	472	166	22,605	117	22,722	10.51	1.34
(%)	(0.1)	(0.3)	(0.7)	(2.7)	(7.3)	(18.6)	(34.0)	(24.9)	(8.7)	(2.1)	(0.7)	(100.0)				
120～	27	64	187	724	2,388	7,336	14,396	11,462	4,045	1,010	375	42,014	187	42,201	10.66	1.27
(%)	(0.1)	(0.2)	(0.4)	(1.7)	(5.7)	(17.5)	(34.3)	(27.3)	(9.6)	(2.4)	(0.9)	(100.0)				
140～	30	65	153	620	2,251	7,787	16,843	13,957	5,029	1,268	434	48,437	222	48,659	10.75	1.23
(%)	(0.1)	(0.1)	(0.3)	(1.3)	(4.6)	(16.1)	(34.8)	(28.8)	(10.4)	(2.6)	(0.9)	(100.0)				
160～	30	38	76	390	1,588	6,040	13,798	12,046	4,460	1,154	413	40,033	210	40,243	10.83	1.21
(%)	(0.1)	(0.1)	(0.2)	(1.0)	(4.0)	(15.1)	(34.5)	(30.1)	(11.1)	(2.9)	(1.0)	(100.0)				
180～	13	29	46	233	893	3,618	8,668	7,712	3,070	808	287	25,377	148	25,525	10.88	1.22
(%)	(0.1)	(0.1)	(0.2)	(0.9)	(3.5)	(14.3)	(34.2)	(30.4)	(12.1)	(3.2)	(1.1)	(100.0)				
200～	7	8	28	90	483	1,943	4,526	4,232	1,667	469	185	13,638	85	13,723	10.92	1.22
(%)	(0.1)	(0.1)	(0.2)	(0.7)	(3.5)	(14.2)	(33.2)	(31.0)	(12.2)	(3.4)	(1.4)	(100.0)				
220～	4	4	14	49	191	852	1,962	1,843	782	229	106	6,036	35	6,071	10.95	1.26
(%)	(0.1)	(0.1)	(0.2)	(0.8)	(3.2)	(14.1)	(32.5)	(30.5)	(13.0)	(3.8)	(1.8)	(100.0)				
240～		2	6	27	83	340	736	738	303	114	36	2,385	15	2,400	10.96	1.28
(%)		(0.1)	(0.3)	(1.1)	(3.5)	(14.3)	(30.9)	(30.9)	(12.7)	(4.8)	(1.5)	(100.0)				
260～	1	1	6	13	77	244	473	436	227	73	22	1,573	13	1,586	10.92	1.33
(%)	(0.1)	(0.1)	(0.4)	(0.8)	(4.9)	(15.5)	(30.1)	(27.7)	(14.4)	(4.6)	(1.4)	(100.0)				
合計	143	302	788	3,139	10,428	33,981	71,442	59,718	22,115	5,742	2,088	209,886	1,074	210,960	10.74	1.27
(%)	(0.1)	(0.1)	(0.4)	(1.5)	(5.0)	(16.2)	(34.0)	(28.5)	(10.5)	(2.7)	(1.0)	(100.0)				
記載なし	160	127	372	1,296	4,126	12,008	22,470	18,455	6,739	1,566	585	67,904	31,244	99,148	10.62	1.33
(%)	(0.2)	(0.2)	(0.5)	(1.9)	(6.1)	(17.7)	(33.1)	(27.2)	(9.9)	(2.3)	(0.9)	(100.0)				
総計	303	429	1,160	4,435	14,554	45,989	93,912	78,173	28,854	7,308	2,673	277,790	32,318	310,108	10.71	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.2)	(16.6)	(33.8)	(28.1)	(10.4)	(2.6)	(1.0)	(100.0)				
平均	148.11	139.99	136.74	139.15	145.92	152.39	155.78	158.28	160.35	162.65	162.86	155.84	159.08	155.86		
標準偏差	39.25	35.54	38.56	36.68	36.63	35.83	34.93	35.06	36.01	37.75	37.87	35.72	38.05	35.74		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査