

表 3629 透析前ヘモグロビン濃度(g/dL) 医療機関別（施設血液透析患者 透析回数週3回）

医療機関	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
国公立大学		1	6	24	60	86	99	91	11	3	2	383	31	414	9.99	1.42
(%)		(0.3)	(1.6)	(6.3)	(15.7)	(22.5)	(25.8)	(23.8)	(2.9)	(0.8)	(0.5)	(100.0)				
私立大学	2	3	12	66	127	279	425	421	156	42	13	1,546	22	1,568	10.49	1.46
(%)	(0.1)	(0.2)	(0.8)	(4.3)	(8.2)	(18.0)	(27.5)	(27.2)	(10.1)	(2.7)	(0.8)	(100.0)				
国立	2		9	17	47	91	132	113	36	14	2	463	32	495	10.35	1.50
(%)	(0.4)		(1.9)	(3.7)	(10.2)	(19.7)	(28.5)	(24.4)	(7.8)	(3.0)	(0.4)	(100.0)				
県市町村立	10	24	83	286	767	2,157	3,937	3,065	1,220	359	124	12,032	798	12,830	10.62	1.36
(%)	(0.1)	(0.2)	(0.7)	(2.4)	(6.4)	(17.9)	(32.7)	(25.5)	(10.1)	(3.0)	(1.0)	(100.0)				
社会保険	3	5	22	76	174	441	831	758	296	83	33	2,722	74	2,796	10.67	1.41
(%)	(0.1)	(0.2)	(0.8)	(2.8)	(6.4)	(16.2)	(30.5)	(27.8)	(10.9)	(3.0)	(1.2)	(100.0)				
厚生連	2	6	36	115	354	1,030	2,118	1,696	777	190	72	6,396	192	6,588	10.75	1.32
(%)	(0.0)	(0.1)	(0.6)	(1.8)	(5.5)	(16.1)	(33.1)	(26.5)	(12.1)	(3.0)	(1.1)	(100.0)				
その他公的	11	41	57	144	444	1,035	2,024	1,926	798	214	92	6,786	479	7,265	10.70	1.45
(%)	(0.2)	(0.6)	(0.8)	(2.1)	(6.5)	(15.3)	(29.8)	(28.4)	(11.8)	(3.2)	(1.4)	(100.0)				
私立総合	1	19	41	99	256	772	1,532	1,401	529	170	52	4,872	112	4,984	10.72	1.38
(%)	(0.0)	(0.4)	(0.8)	(2.0)	(5.3)	(15.8)	(31.4)	(28.8)	(10.9)	(3.5)	(1.1)	(100.0)				
私立	73	132	356	1,345	4,136	11,032	20,586	17,026	6,669	1,778	612	63,745	2,854	66,599	10.65	1.35
(%)	(0.1)	(0.2)	(0.6)	(2.1)	(6.5)	(17.3)	(32.3)	(26.7)	(10.5)	(2.8)	(1.0)	(100.0)				
私立診療所	157	131	320	1,406	5,361	20,115	43,533	34,982	11,878	2,684	931	121,498	3,934	125,432	10.72	1.20
(%)	(0.1)	(0.1)	(0.3)	(1.2)	(4.4)	(16.6)	(35.8)	(28.8)	(9.8)	(2.2)	(0.8)	(100.0)				
総計	261	362	942	3,578	11,726	37,038	75,217	61,479	22,370	5,537	1,933	220,443	8,528	228,971	10.69	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.3)	(16.8)	(34.1)	(27.9)	(10.1)	(2.5)	(0.9)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査