

表 3669 総コレステロール濃度(mg/dL) 週透析回数(回/週)別 (透析患者全体)

週透析回数	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	合計	記載なし	総計	平均	標準偏差
1回	25	83	210	310	277	201	132	52	21	18	1,329	511	1,840	165.25	35.95
(%)	(1.9)	(6.2)	(15.8)	(23.3)	(20.8)	(15.1)	(9.9)	(3.9)	(1.6)	(1.4)	(100.0)				
2回	163	453	813	951	823	536	332	158	66	59	4,354	1,838	6,192	158.85	38.63
(%)	(3.7)	(10.4)	(18.7)	(21.8)	(18.9)	(12.3)	(7.6)	(3.6)	(1.5)	(1.4)	(100.0)				
3回	7,483	21,733	40,292	46,056	37,867	23,663	12,564	5,475	2,131	1,334	198,598	72,041	270,639	155.24	35.36
(%)	(3.8)	(10.9)	(20.3)	(23.2)	(19.1)	(11.9)	(6.3)	(2.8)	(1.1)	(0.7)	(100.0)				
4回	33	65	110	143	155	117	67	27	16	11	744	250	994	163.46	38.97
(%)	(4.4)	(8.7)	(14.8)	(19.2)	(20.8)	(15.7)	(9.0)	(3.6)	(2.2)	(1.5)	(100.0)				
5回	1	3	6	8	16	12	8	4	7	2	67	22	89	183.48	42.96
(%)	(1.5)	(4.5)	(9.0)	(11.9)	(23.9)	(17.9)	(11.9)	(6.0)	(10.4)	(3.0)	(100.0)				
6回		4	3	9	8	7	9	3		1	44	37	81	175.45	37.05
(%)		(9.1)	(6.8)	(20.5)	(18.2)	(15.9)	(20.5)	(6.8)		(2.3)	(100.0)				
7回		1	2	3	3	5	3	4	1	2	24	6	30	195.13	48.58
(%)		(4.2)	(8.3)	(12.5)	(12.5)	(20.8)	(12.5)	(16.7)	(4.2)	(8.3)	(100.0)				
合計	7,705	22,342	41,436	47,480	39,149	24,541	13,115	5,723	2,242	1,427	205,160	74,705	279,865	155.43	35.48
(%)	(3.8)	(10.9)	(20.2)	(23.1)	(19.1)	(12.0)	(6.4)	(2.8)	(1.1)	(0.7)	(100.0)				
記載なし	125	380	765	1,179	1,094	984	608	348	158	159	5,800	24,443	30,243	171.14	41.08
(%)	(2.2)	(6.6)	(13.2)	(20.3)	(18.9)	(17.0)	(10.5)	(6.0)	(2.7)	(2.7)	(100.0)				
総計	7,830	22,722	42,201	48,659	40,243	25,525	13,723	6,071	2,400	1,586	210,960	99,148	310,108	155.86	35.74
(%)	(3.7)	(10.8)	(20.0)	(23.1)	(19.1)	(12.1)	(6.5)	(2.9)	(1.1)	(0.8)	(100.0)				
平均	2.98	2.98	2.97	2.97	2.97	2.97	2.96	2.96	2.97	2.95	2.97	2.97	2.97		
標準偏差	0.20	0.20	0.21	0.23	0.24	0.26	0.29	0.29	0.30	0.36	0.23	0.25	0.24		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査