

表 4024 APD使用の有無 原疾患別 (PD・HD(F)等併用患者)

原疾患	なし	あり	合計	不明	記載なし	総計
慢性糸球体腎炎	235	222	457		270	727
(%)	(51.4)	(48.6)	(100.0)			
慢性腎盂腎炎	6	5	11		3	14
(%)	(54.5)	(45.5)	(100.0)			
急速進行性糸球体腎炎	4	2	6		4	10
(%)	(66.7)	(33.3)	(100.0)			
妊娠腎/妊娠中毒症	5	2	7		3	10
(%)	(71.4)	(28.6)	(100.0)			
その他の分類不能の腎炎	4	7	11		8	19
(%)	(36.4)	(63.6)	(100.0)			
多発性嚢胞腎	11	12	23	1	13	37
(%)	(47.8)	(52.2)	(100.0)			
腎硬化症	39	40	79	1	63	143
(%)	(49.4)	(50.6)	(100.0)			
悪性高血圧	6	9	15		7	22
(%)	(40.0)	(60.0)	(100.0)			
糖尿病性腎症	141	143	284	2	244	530
(%)	(49.6)	(50.4)	(100.0)			
SLE腎炎	2	3	5		3	8
(%)	(40.0)	(60.0)	(100.0)			
アミロイド腎					2	2
(%)						
痛風腎	5	1	6		1	7
(%)	(83.3)	(16.7)	(100.0)			
先天性代謝異常による腎不全		1	1			1
(%)		(100.0)	(100.0)			
腎・尿路結核						
(%)						
腎・尿路結石		1	1		1	2
(%)		(100.0)	(100.0)			
腎・尿路腫瘍		2	2		1	3
(%)		(100.0)	(100.0)			
閉塞性尿路障害		1	1		2	3
(%)		(100.0)	(100.0)			
骨髄腫					1	1
(%)						
腎形成不全	1	2	3		1	4
(%)	(33.3)	(66.7)	(100.0)			
不明	47	28	75	1	74	150
(%)	(62.7)	(37.3)	(100.0)			
再導入	3	6	9		6	15
(%)	(33.3)	(66.7)	(100.0)			
その他	9	12	21		19	40
(%)	(42.9)	(57.1)	(100.0)			
合計	518	499	1,017	5	726	1,748
(%)	(50.9)	(49.1)	(100.0)			
記載なし						
(%)						
総計	518	499	1,017	5	726	1,748
(%)	(50.9)	(49.1)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査