

表 4160 PET Cr D/P比 原疾患別（腹膜透析単独患者）

原疾患	0.5未満	0.5～	0.65～	0.81～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	101	340	342	108	891	1,438	2,329	0.65	0.14
(%)	(11.3)	(38.2)	(38.4)	(12.1)	(100.0)				
慢性腎盂腎炎	6	12	9	5	32	47	79	0.64	0.15
(%)	(18.8)	(37.5)	(28.1)	(15.6)	(100.0)				
急速進行性糸球体腎炎	1	8	13	2	24	40	64	0.68	0.10
(%)	(4.2)	(33.3)	(54.2)	(8.3)	(100.0)				
妊娠腎/妊娠中毒症	1	8	3	1	13	9	22	0.61	0.11
(%)	(7.7)	(61.5)	(23.1)	(7.7)	(100.0)				
その他の分類不能の腎炎	5	8	10	2	25	41	66	0.62	0.12
(%)	(20.0)	(32.0)	(40.0)	(8.0)	(100.0)				
多発性嚢胞腎	8	33	24	7	72	134	206	0.63	0.12
(%)	(11.1)	(45.8)	(33.3)	(9.7)	(100.0)				
腎硬化症	20	97	151	38	306	634	940	0.67	0.12
(%)	(6.5)	(31.7)	(49.3)	(12.4)	(100.0)				
悪性高血圧	6	8	8	1	23	62	85	0.59	0.12
(%)	(26.1)	(34.8)	(34.8)	(4.3)	(100.0)				
糖尿病性腎症	46	235	328	123	732	1,503	2,235	0.68	0.14
(%)	(6.3)	(32.1)	(44.8)	(16.8)	(100.0)				
SLE腎炎	3	6	7	1	17	37	54	0.61	0.13
(%)	(17.6)	(35.3)	(41.2)	(5.9)	(100.0)				
アミロイド腎		1	1		2	5	7	0.70	0.12
(%)		(50.0)	(50.0)		(100.0)				
痛風腎		4	5	1	10	16	26	0.66	0.09
(%)		(40.0)	(50.0)	(10.0)	(100.0)				
先天性代謝異常による腎不全	2	1	1	2	6	11	17	0.66	0.23
(%)	(33.3)	(16.7)	(16.7)	(33.3)	(100.0)				
腎・尿路結核		1			1		1	0.53	
(%)		(100.0)			(100.0)				
腎・尿路結石		4		1	5	9	14	0.66	0.13
(%)		(80.0)		(20.0)	(100.0)				
腎・尿路腫瘍	1		1	2	4	7	11	0.71	0.17
(%)	(25.0)		(25.0)	(50.0)	(100.0)				
閉塞性尿路障害	1	4	2	2	9	10	19	0.65	0.13
(%)	(11.1)	(44.4)	(22.2)	(22.2)	(100.0)				
骨髄腫		1			1	6	7	0.60	
(%)		(100.0)			(100.0)				
腎形成不全	4	7	13		24	48	72	0.62	0.12
(%)	(16.7)	(29.2)	(54.2)		(100.0)				
不明	20	92	85	26	223	466	689	0.65	0.13
(%)	(9.0)	(41.3)	(38.1)	(11.7)	(100.0)				
再導入	3	9	6	1	19	17	36	0.62	0.11
(%)	(15.8)	(47.4)	(31.6)	(5.3)	(100.0)				
その他	11	21	30	12	74	135	209	0.67	0.14
(%)	(14.9)	(28.4)	(40.5)	(16.2)	(100.0)				
合計	239	900	1,039	335	2,513	4,675	7,188	0.66	0.14
(%)	(9.5)	(35.8)	(41.3)	(13.3)	(100.0)				
記載なし									
(%)									
総計	239	900	1,039	335	2,513	4,675	7,188	0.66	0.14
(%)	(9.5)	(35.8)	(41.3)	(13.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査