

表 4230 残腎Kt/V Non-HDL-C濃度(mg/dL)別（腹膜透析患者）

Non-HDL-C濃度	0.1未満	0.1～	0.4～	0.8～	1.2～	1.7～	2.0～	2.4～	合計	記載なし	総計	平均	標準偏差
50未満	2	2	4	2	1				11	7	18	0.51	0.44
(%)	(18.2)	(18.2)	(36.4)	(18.2)	(9.1)				(100.0)				
50～	18	13	8	3	4			2	48	66	114	0.44	0.68
(%)	(37.5)	(27.1)	(16.7)	(6.3)	(8.3)			(4.2)	(100.0)				
70～	42	24	31	19	12	3	2	1	134	217	351	0.52	0.69
(%)	(31.3)	(17.9)	(23.1)	(14.2)	(9.0)	(2.2)	(1.5)	(0.7)	(100.0)				
90～	78	61	42	31	27	2	4	3	248	412	660	0.53	0.79
(%)	(31.5)	(24.6)	(16.9)	(12.5)	(10.9)	(0.8)	(1.6)	(1.2)	(100.0)				
110～	86	62	75	51	17	7	4	5	307	490	797	0.55	0.71
(%)	(28.0)	(20.2)	(24.4)	(16.6)	(5.5)	(2.3)	(1.3)	(1.6)	(100.0)				
130～	60	60	71	47	16	7	5	2	268	394	662	0.54	0.54
(%)	(22.4)	(22.4)	(26.5)	(17.5)	(6.0)	(2.6)	(1.9)	(0.7)	(100.0)				
150～	36	37	40	26	17	4		1	161	275	436	0.54	0.52
(%)	(22.4)	(23.0)	(24.8)	(16.1)	(10.6)	(2.5)		(0.6)	(100.0)				
170～	33	20	20	16	7	4	2	1	103	160	263	0.51	0.61
(%)	(32.0)	(19.4)	(19.4)	(15.5)	(6.8)	(3.9)	(1.9)	(1.0)	(100.0)				
190～	6	3	15	5	5	1		3	38	82	120	0.88	1.05
(%)	(15.8)	(7.9)	(39.5)	(13.2)	(13.2)	(2.6)		(7.9)	(100.0)				
210～	9	4	9	3		1	3	2	31	59	90	0.70	0.85
(%)	(29.0)	(12.9)	(29.0)	(9.7)		(3.2)	(9.7)	(6.5)	(100.0)				
合計	370	286	315	203	106	29	20	20	1,349	2,162	3,511	0.55	0.68
(%)	(27.4)	(21.2)	(23.4)	(15.0)	(7.9)	(2.1)	(1.5)	(1.5)	(100.0)				
記載なし	140	183	194	146	89	27	19	9	807	4,623	5,430	0.64	0.62
(%)	(17.3)	(22.7)	(24.0)	(18.1)	(11.0)	(3.3)	(2.4)	(1.1)	(100.0)				
総計	510	469	509	349	195	56	39	29	2,156	6,785	8,941	0.58	0.66
(%)	(23.7)	(21.8)	(23.6)	(16.2)	(9.0)	(2.6)	(1.8)	(1.3)	(100.0)				
平均	123.81	124.80	131.64	129.05	124.51	140.38	137.35	141.05	127.51	129.07	128.47		
標準偏差	38.75	34.32	40.50	33.25	36.30	37.21	47.58	54.78	37.84	38.12	38.01		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査