

表 4334 PD Kt/V 腹膜炎罹患回数(回/年)別 (腹膜透析患者)

腹膜炎罹患回数	0.1未満	0.1～	0.4～	0.8～	1.2～	1.7～	2.0～	2.4～	合計	記載なし	総計	平均	標準偏差
なし	16	133	220	421	753	282	142	82	2,049	2,602	4,651	1.32	0.78
(%)	(0.8)	(6.5)	(10.7)	(20.5)	(36.7)	(13.8)	(6.9)	(4.0)	(100.0)				
1回	2	22	24	48	70	23	14	8	211	316	527	1.23	0.69
(%)	(0.9)	(10.4)	(11.4)	(22.7)	(33.2)	(10.9)	(6.6)	(3.8)	(100.0)				
2回	1	2	5	3	14	4	4		33	49	82	1.26	0.58
(%)	(3.0)	(6.1)	(15.2)	(9.1)	(42.4)	(12.1)	(12.1)		(100.0)				
3回		2		1	2		1	1	7	20	27	1.64	1.84
(%)		(28.6)		(14.3)	(28.6)		(14.3)	(14.3)	(100.0)				
4回			3		2				5	2	7	0.92	0.40
(%)			(60.0)		(40.0)				(100.0)				
5回				1	1	1			3	1	4	1.37	0.50
(%)				(33.3)	(33.3)	(33.3)			(100.0)				
6回										1	1		
(%)													
7回													
(%)													
8回										1	1		
(%)													
9回以上										1	1		
(%)													
合計	19	159	252	474	842	310	161	91	2,308	2,993	5,301	1.31	0.77
(%)	(0.8)	(6.9)	(10.9)	(20.5)	(36.5)	(13.4)	(7.0)	(3.9)	(100.0)				
不明			15	15	13	2			45	34	79	1.02	0.35
(%)			(33.3)	(33.3)	(28.9)	(4.4)			(100.0)				
記載なし		4	2	9	11	4	2		32	3,529	3,561	1.16	0.54
(%)		(12.5)	(6.3)	(28.1)	(34.4)	(12.5)	(6.3)		(100.0)				
総計	19	163	269	498	866	316	163	91	2,385	6,556	8,941	1.30	0.76
(%)	(0.8)	(6.8)	(11.3)	(20.9)	(36.3)	(13.2)	(6.8)	(3.8)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査