

表 4355 PD Kt/V 出口部感染罹患回数(回/年)別 (腹膜透析単独患者)

| 出口部感染<br>罹患回数 | 0.1未満 | 0.1～   | 0.4～   | 0.8～   | 1.2～   | 1.7～   | 2.0～   | 2.4～  | 合計      | 記載なし  | 総計    | 平均   | 標準偏差 |
|---------------|-------|--------|--------|--------|--------|--------|--------|-------|---------|-------|-------|------|------|
| なし            | 8     | 62     | 154    | 311    | 519    | 211    | 109    | 71    | 1,445   | 2,020 | 3,465 | 1.37 | 0.81 |
| (%)           | (0.6) | (4.3)  | (10.7) | (21.5) | (35.9) | (14.6) | (7.5)  | (4.9) | (100.0) |       |       |      |      |
| 1回            | 2     | 41     | 40     | 65     | 97     | 24     | 19     | 7     | 295     | 270   | 565   | 1.12 | 0.62 |
| (%)           | (0.7) | (13.9) | (13.6) | (22.0) | (32.9) | (8.1)  | (6.4)  | (2.4) | (100.0) |       |       |      |      |
| 2回            |       | 2      | 6      | 21     | 32     | 8      | 7      | 1     | 77      | 73    | 150   | 1.28 | 0.49 |
| (%)           |       | (2.6)  | (7.8)  | (27.3) | (41.6) | (10.4) | (9.1)  | (1.3) | (100.0) |       |       |      |      |
| 3回            |       |        | 5      | 5      | 14     | 1      | 2      | 1     | 28      | 17    | 45    | 1.41 | 1.04 |
| (%)           |       |        | (17.9) | (17.9) | (50.0) | (3.6)  | (7.1)  | (3.6) | (100.0) |       |       |      |      |
| 4回            |       | 4      | 3      | 5      | 5      |        |        |       | 17      | 7     | 24    | 0.82 | 0.47 |
| (%)           |       | (23.5) | (17.6) | (29.4) | (29.4) |        |        |       | (100.0) |       |       |      |      |
| 5回            |       |        | 3      |        | 2      | 1      | 1      |       | 7       | 6     | 13    | 1.20 | 0.57 |
| (%)           |       |        | (42.9) |        | (28.6) | (14.3) | (14.3) |       | (100.0) |       |       |      |      |
| 6回            |       |        |        | 2      | 2      | 1      |        |       | 5       | 7     | 12    | 1.30 | 0.33 |
| (%)           |       |        |        | (40.0) | (40.0) | (20.0) |        |       | (100.0) |       |       |      |      |
| 7回            |       |        |        |        |        |        |        |       |         |       |       |      |      |
| (%)           |       |        |        |        |        |        |        |       |         |       |       |      |      |
| 8回            |       |        |        |        |        |        |        |       |         | 1     | 1     |      |      |
| (%)           |       |        |        |        |        |        |        |       |         |       |       |      |      |
| 9回以上          |       | 2      |        | 1      | 4      |        |        |       | 7       | 7     | 14    | 1.00 | 0.56 |
| (%)           |       | (28.6) |        | (14.3) | (57.1) |        |        |       | (100.0) |       |       |      |      |
| 合計            | 10    | 111    | 211    | 410    | 675    | 246    | 138    | 80    | 1,881   | 2,408 | 4,289 | 1.32 | 0.78 |
| (%)           | (0.5) | (5.9)  | (11.2) | (21.8) | (35.9) | (13.1) | (7.3)  | (4.3) | (100.0) |       |       |      |      |
| 不明            |       |        | 13     | 10     | 8      | 1      |        |       | 32      | 29    | 61    | 0.97 | 0.36 |
| (%)           |       |        | (40.6) | (31.3) | (25.0) | (3.1)  |        |       | (100.0) |       |       |      |      |
| 記載なし          |       | 5      | 2      | 9      | 10     | 2      | 2      |       | 30      | 2,808 | 2,838 | 1.08 | 0.56 |
| (%)           |       | (16.7) | (6.7)  | (30.0) | (33.3) | (6.7)  | (6.7)  |       | (100.0) |       |       |      |      |
| 総計            | 10    | 116    | 226    | 429    | 693    | 249    | 140    | 80    | 1,943   | 5,245 | 7,188 | 1.31 | 0.77 |
| (%)           | (0.5) | (6.0)  | (11.6) | (22.1) | (35.7) | (12.8) | (7.2)  | (4.1) | (100.0) |       |       |      |      |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査