

表 4458 腹膜炎罹患回数(回/年) 使用透析液の種類別(腹膜透析患者)

使用透析液の種類	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
1.5%液のみ	1,893	199	34	9	1	1					2,137	11	127	2,275
(%)	(88.6)	(9.3)	(1.6)	(0.4)	(0.0)	(0.0)					(100.0)			
1.5%液と2.5%液の併用	624	83	14	6	1	1					729	9	49	787
(%)	(85.6)	(11.4)	(1.9)	(0.8)	(0.1)	(0.1)					(100.0)			
2.5%液のみ	138	18	6	1							163	2	12	177
(%)	(84.7)	(11.0)	(3.7)	(0.6)							(100.0)			
4.25%液のみ(イコデキストリンなし)	5	3									8			8
(%)	(62.5)	(37.5)									(100.0)			
イコデキストリン単独(ブドウ糖使用なし)	53	2									55		2	57
(%)	(96.4)	(3.6)									(100.0)			
1.5%+イコデキストリン	1,009	119	13	3	2	1	1		1	1	1,150	2	70	1,222
(%)	(87.7)	(10.3)	(1.1)	(0.3)	(0.2)	(0.1)	(0.1)		(0.1)	(0.1)	(100.0)			
1.5%+2.5%+イコデキストリン	476	40	10	3	2						531		28	559
(%)	(89.6)	(7.5)	(1.9)	(0.6)	(0.4)						(100.0)			
2.5%+イコデキストリン	358	46	5	5	1	1					416	3	22	441
(%)	(86.1)	(11.1)	(1.2)	(1.2)	(0.2)	(0.2)					(100.0)			
4.25%液+イコデキストリン	5	1									6			6
(%)	(83.3)	(16.7)									(100.0)			
合計	4,561	511	82	27	7	4	1		1	1	5,195	27	310	5,532
(%)	(87.8)	(9.8)	(1.6)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			
不明	11										11		52	2
(%)	(100.0)										(100.0)			65
記載なし	79	16									95		3,249	3,344
(%)	(83.2)	(16.8)									(100.0)			
総計	4,651	527	82	27	7	4	1		1	1	5,301	79	3,561	8,941
(%)	(87.7)	(9.9)	(1.5)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)	(0.0)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査