

表 4531 出口部感染罹患回数(回/年) PTH値(pg/mL)別 (腹膜透析患者) PTH測定法:intact-PTH

| PTH値 | なし | 1回 | 2回 | 3回 | 4回 | 5回 | 6回 | 7回 | 8回 | 9回以上 | 合計 | 不明 | 記載なし | 総計 |
|------|--------|--------|--------|--------|--------|--------|--------|-------|-------|--------|---------|--------|--------|--------|
| 20未満 | 75 | 8 | 5 | | | 1 | | | 1 | 1 | 91 | | 32 | 123 |
| (%) | (82.4) | (8.8) | (5.5) | | | (1.1) | | | (1.1) | (1.1) | (100.0) | | | |
| 20～ | 161 | 35 | 7 | 1 | | | 2 | | 1 | 1 | 208 | 3 | 70 | 281 |
| (%) | (77.4) | (16.8) | (3.4) | (0.5) | | | (1.0) | | (0.5) | (0.5) | (100.0) | | | |
| 40～ | 177 | 25 | 3 | 2 | | | 1 | | | | 208 | | 3 | 268 |
| (%) | (85.1) | (12.0) | (1.4) | (1.0) | | | (0.5) | | | | (100.0) | | | |
| 60～ | 187 | 41 | 9 | 2 | 1 | | 6 | | | 1 | 247 | 1 | 67 | 315 |
| (%) | (75.7) | (16.6) | (3.6) | (0.8) | (0.4) | | (2.4) | | | (0.4) | (100.0) | | | |
| 80～ | 217 | 30 | 11 | 2 | 1 | | 2 | 2 | | 1 | 266 | 2 | 41 | 309 |
| (%) | (81.6) | (11.3) | (4.1) | (0.8) | (0.4) | | (0.8) | (0.8) | | (0.4) | (100.0) | | | |
| 100～ | 212 | 34 | 8 | | 1 | 1 | | | | 2 | 258 | 1 | 57 | 316 |
| (%) | (82.2) | (13.2) | (3.1) | | (0.4) | (0.4) | | | | (0.8) | (100.0) | | | |
| 120～ | 207 | 28 | 6 | | 2 | | 1 | | | 1 | 245 | 1 | 59 | 305 |
| (%) | (84.5) | (11.4) | (2.4) | | (0.8) | | (0.4) | | | (0.4) | (100.0) | | | |
| 140～ | 184 | 31 | 6 | 3 | | 4 | | | 1 | 1 | 230 | 1 | 54 | 285 |
| (%) | (80.0) | (13.5) | (2.6) | (1.3) | | (1.7) | | | (0.4) | (0.4) | (100.0) | | | |
| 160～ | 174 | 30 | 9 | 4 | 2 | | | | | | 219 | 2 | 55 | 276 |
| (%) | (79.5) | (13.7) | (4.1) | (1.8) | (0.9) | | | | | | (100.0) | | | |
| 180～ | 454 | 50 | 12 | 9 | 1 | | 1 | | | 2 | 529 | 2 | 123 | 654 |
| (%) | (85.8) | (9.5) | (2.3) | (1.7) | (0.2) | | (0.2) | | | (0.4) | (100.0) | | | |
| 240～ | 7 | 1 | 1 | | | | | | | | 9 | | 1 | 10 |
| (%) | (77.8) | (11.1) | (11.1) | | | | | | | | (100.0) | | | |
| 241～ | 280 | 45 | 12 | 3 | 2 | 2 | | | | | 344 | 3 | 84 | 431 |
| (%) | (81.4) | (13.1) | (3.5) | (0.9) | (0.6) | (0.6) | | | | | (100.0) | | | |
| 300～ | 238 | 33 | 11 | 3 | 2 | 1 | 1 | | | | 289 | | 42 | 331 |
| (%) | (82.4) | (11.4) | (3.8) | (1.0) | (0.7) | (0.3) | (0.3) | | | | (100.0) | | | |
| 360～ | 121 | 25 | 6 | 3 | 1 | 2 | | | | | 158 | 2 | 40 | 200 |
| (%) | (76.6) | (15.8) | (3.8) | (1.9) | (0.6) | (1.3) | | | | | (100.0) | | | |
| 420～ | 91 | 9 | 5 | 2 | | 1 | 1 | | | 1 | 110 | | 29 | 139 |
| (%) | (82.7) | (8.2) | (4.5) | (1.8) | | (0.9) | (0.9) | | | (0.9) | (100.0) | | | |
| 480～ | 27 | 4 | | 3 | | | | | | | 34 | | 9 | 43 |
| (%) | (79.4) | (11.8) | | (8.8) | | | | | | | (100.0) | | | |
| 500～ | 32 | 5 | | 1 | 2 | | | | | 1 | 41 | 1 | 12 | 54 |
| (%) | (78.0) | (12.2) | | (2.4) | (4.9) | | | | | (2.4) | (100.0) | | | |
| 540～ | 45 | 7 | 1 | 1 | 1 | | 1 | | | | 56 | | 17 | 73 |
| (%) | (80.4) | (12.5) | (1.8) | (1.8) | (1.8) | | (1.8) | | | | (100.0) | | | |
| 600～ | 66 | 16 | 3 | 1 | | | | | | | 86 | | 19 | 105 |
| (%) | (76.7) | (18.6) | (3.5) | (1.2) | | | | | | | (100.0) | | | |
| 800～ | 71 | 6 | 2 | | | | | | | | 79 | | 31 | 110 |
| (%) | (89.9) | (7.6) | (2.5) | | | | | | | | (100.0) | | | |
| 合計 | 3,026 | 463 | 117 | 40 | 16 | 20 | 10 | | 3 | 12 | 3,707 | 22 | 899 | 4,628 |
| (%) | (81.6) | (12.5) | (3.2) | (1.1) | (0.4) | (0.5) | (0.3) | | (0.1) | (0.3) | (100.0) | | | |
| 記載なし | 221 | 28 | 11 | 2 | 5 | | 1 | | | 1 | 269 | 2 | 105 | 376 |
| (%) | (82.2) | (10.4) | (4.1) | (0.7) | (1.9) | | (0.4) | | | (0.4) | (100.0) | | | |
| 総計 | 3,247 | 491 | 128 | 42 | 21 | 20 | 11 | | 3 | 13 | 3,976 | 24 | 1,004 | 5,004 |
| (%) | (81.7) | (12.3) | (3.2) | (1.1) | (0.5) | (0.5) | (0.3) | | (0.1) | (0.3) | (100.0) | | | |
| 平均 | 223.79 | 213.06 | 209.44 | 269.07 | 265.56 | 167.90 | 195.80 | | 64.67 | 171.33 | 221.99 | 170.55 | 232.97 | 223.88 |
| 標準偏差 | 218.90 | 188.66 | 169.98 | 165.95 | 163.22 | 129.30 | 188.89 | | 72.76 | 153.89 | 212.59 | 137.61 | 289.51 | 229.31 |

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査