

表 4532 出口部感染罹患回数(回/年) PTH値(pg/mL)別(腹膜透析患者) PTH測定法:whole-PTH

PTH値	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
20未満	27	6	3		1						37		3	40
(%)	(73.0)	(16.2)	(8.1)		(2.7)						(100.0)			
20～	31	7	3	3	1						45			45
(%)	(68.9)	(15.6)	(6.7)	(6.7)	(2.2)						(100.0)			
35～	41	5	3				1				50		5	55
(%)	(82.0)	(10.0)	(6.0)				(2.0)				(100.0)			
50～	52	13	1	2	1		2				71		6	77
(%)	(73.2)	(18.3)	(1.4)	(2.8)	(1.4)		(2.8)				(100.0)			
75～	54	9	1	1		1					66		3	69
(%)	(81.8)	(13.6)	(1.5)	(1.5)		(1.5)					(100.0)			
100～	52	13	2	1							68		3	71
(%)	(76.5)	(19.1)	(2.9)	(1.5)							(100.0)			
125～	34	3	3								40		5	45
(%)	(85.0)	(7.5)	(7.5)								(100.0)			
150～	1										1			1
(%)	(100.0)										(100.0)			
151～	32	7									39		4	43
(%)	(82.1)	(17.9)									(100.0)			
180～	22	4	1	1						1	29		5	34
(%)	(75.9)	(13.8)	(3.4)	(3.4)						(3.4)	(100.0)			
210～	17	5	5								27			27
(%)	(63.0)	(18.5)	(18.5)								(100.0)			
240～	17	2	1	1							21		2	23
(%)	(81.0)	(9.5)	(4.8)	(4.8)							(100.0)			
300～	17	1									18		1	19
(%)	(94.4)	(5.6)									(100.0)			
350～	7		1								8			8
(%)	(87.5)		(12.5)								(100.0)			
400～	7	2	1	1							11			11
(%)	(63.6)	(18.2)	(9.1)	(9.1)							(100.0)			
500～	10			1							11			11
(%)	(90.9)			(9.1)							(100.0)			
合計	421	77	25	11	3	1	3			1	542		37	579
(%)	(77.7)	(14.2)	(4.6)	(2.0)	(0.6)	(0.2)	(0.6)			(0.2)	(100.0)			
記載なし	69	42	1								112		28	140
(%)	(61.6)	(37.5)	(0.9)								(100.0)			
総計	490	119	26	11	3	1	3			1	654		65	719
(%)	(74.9)	(18.2)	(4.0)	(1.7)	(0.5)	(0.2)	(0.5)			(0.2)	(100.0)			
平均	137.35	111.55	136.20	174.55	36.33	96.00	61.00			208.00	133.46		119.30	132.56
標準偏差	138.00	88.66	116.11	203.99	23.18		19.16				132.17		78.12	129.39

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査