

表 212 2015年末患者, 性別, 原疾患別

原疾患	男性	女性	合計	記載なし	総計
慢性糸球体腎炎	56,150	37,197	93,347		93,347
(%)	(27.9)	(33.2)	(29.8)		(29.8)
慢性腎盂腎炎	1,362	1,573	2,935		2,935
(%)	(0.7)	(1.4)	(0.9)		(0.9)
急速進行性糸球体腎炎	1,287	1,191	2,478		2,478
(%)	(0.6)	(1.1)	(0.8)		(0.8)
妊娠腎/妊娠中毒症	12	1,533	1,545		1,545
(%)	(0.0)	(1.4)	(0.5)		(0.5)
その他の分類不能の腎炎	821	584	1,405		1,405
(%)	(0.4)	(0.5)	(0.4)		(0.4)
多発性嚢胞腎	6,183	5,073	11,256		11,256
(%)	(3.1)	(4.5)	(3.6)		(3.6)
腎硬化症	19,745	10,060	29,805		29,805
(%)	(9.8)	(9.0)	(9.5)		(9.5)
悪性高血圧	1,771	847	2,618		2,618
(%)	(0.9)	(0.8)	(0.8)		(0.8)
糖尿病性腎症	84,807	35,471	120,278		120,278
(%)	(42.1)	(31.7)	(38.4)		(38.4)
SLE腎炎	695	1,532	2,227		2,227
(%)	(0.3)	(1.4)	(0.7)		(0.7)
アミロイド腎	200	262	462		462
(%)	(0.1)	(0.2)	(0.1)		(0.1)
痛風腎	939	136	1,075		1,075
(%)	(0.5)	(0.1)	(0.3)		(0.3)
先天性代謝異常による腎不全	159	115	274		274
(%)	(0.1)	(0.1)	(0.1)		(0.1)
腎・尿路結核	106	89	195		195
(%)	(0.1)	(0.1)	(0.1)		(0.1)
腎・尿路結石	379	202	581		581
(%)	(0.2)	(0.2)	(0.2)		(0.2)
腎・尿路腫瘍	701	209	910		910
(%)	(0.3)	(0.2)	(0.3)		(0.3)
閉塞性尿路障害	482	248	730		730
(%)	(0.2)	(0.2)	(0.2)		(0.2)
骨髄腫	184	107	291		291
(%)	(0.1)	(0.1)	(0.1)		(0.1)
腎形成不全	411	249	660		660
(%)	(0.2)	(0.2)	(0.2)		(0.2)
不明	18,564	11,333	29,897		29,897
(%)	(9.2)	(10.1)	(9.5)		(9.5)
再導入	1,434	749	2,183		2,183
(%)	(0.7)	(0.7)	(0.7)		(0.7)
その他	4,932	3,112	8,044		8,044
(%)	(2.4)	(2.8)	(2.6)		(2.6)
合計	201,324	111,872	313,196		313,196
(%)	(100.0)	(100.0)	(100.0)		(100.0)
記載なし	13	8	21		21
総計	201,337	111,880	313,217		313,217

数値下のカッコ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査