

表 478 HDF希釈方法 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
施設血液透析	1				1		248,724	248,725
(%)	(0.0)				(0.0)		(92.8)	(79.4)
Off-line HDF	499	3,772		7	4,278	1	1,053	5,332
(%)	(1.3)	(66.2)		(0.4)	(9.5)	(50.0)	(0.4)	(1.7)
On-line HDF	36,778	1,657	1	317	38,753		5,774	44,527
(%)	(97.3)	(29.1)	(2.2)	(18.7)	(85.7)		(2.2)	(14.2)
Push/Pull HDF	6	29	1	44	80		30	110
(%)	(0.0)	(0.5)	(2.2)	(2.6)	(0.2)		(0.0)	(0.0)
ハイオフィльтраーション	4	133		1	138		129	267
(%)	(0.0)	(2.3)		(0.1)	(0.3)		(0.0)	(0.1)
IHDF	383	67	43	1,327	1,820	1	1,719	3,540
(%)	(1.0)	(1.2)	(95.6)	(78.2)	(4.0)	(50.0)	(0.6)	(1.1)
血液濾過							16	16
(%)							(0.0)	(0.0)
血液吸着透析 (リクセル等使用)							1,376	1,376
(%)							(0.5)	(0.4)
在宅血液透析							555	555
(%)							(0.2)	(0.2)
腹膜透析 (CAPD)	44	13			57		4,845	4,902
(%)	(0.1)	(0.2)			(0.1)		(1.8)	(1.6)
腹膜透析 (APD)	51	14		1	66		2,709	2,775
(%)	(0.1)	(0.2)		(0.1)	(0.1)		(1.0)	(0.9)
腹膜透析 (CCPD)	25	10			35		1,057	1,092
(%)	(0.1)	(0.2)			(0.1)		(0.4)	(0.3)
合計	37,791	5,695	45	1,697	45,228	2	267,987	313,217
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)
記載なし								
総計	37,791	5,695	45	1,697	45,228	2	267,987	313,217

数値下のかっこ内は列方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2015年調査