

表 1810 週透析回数(回/週) 原疾患別 (透析患者全体)

原疾患	1回	2回	3回	4回	5回	6回	7回	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	643	1,440	81,956	350	43	64	12	84,508	8,839	93,347	2.98	0.25
(%)	(0.8)	(1.7)	(97.0)	(0.4)	(0.1)	(0.1)	(0.0)	(100.0)				
慢性腎盂腎炎	22	76	2,549	12	3	1		2,663	272	2,935	2.96	0.27
(%)	(0.8)	(2.9)	(95.7)	(0.5)	(0.1)	(0.0)		(100.0)				
急速進行性糸球体腎炎	16	74	2,128	6	1			2,225	253	2,478	2.96	0.25
(%)	(0.7)	(3.3)	(95.6)	(0.3)	(0.0)			(100.0)				
妊娠腎/妊娠中毒症	6	13	1,421	9				1,449	96	1,545	2.99	0.18
(%)	(0.4)	(0.9)	(98.1)	(0.6)				(100.0)				
その他の分類不能の腎炎	16	39	1,204	3	1	1	1	1,265	140	1,405	2.95	0.32
(%)	(1.3)	(3.1)	(95.2)	(0.2)	(0.1)	(0.1)	(0.1)	(100.0)				
多発性嚢胞腎	48	239	9,944	42	5	4	3	10,285	971	11,256	2.98	0.24
(%)	(0.5)	(2.3)	(96.7)	(0.4)	(0.0)	(0.0)	(0.0)	(100.0)				
腎硬化症	185	742	25,852	63	5	9	1	26,857	2,948	29,805	2.96	0.25
(%)	(0.7)	(2.8)	(96.3)	(0.2)	(0.0)	(0.0)	(0.0)	(100.0)				
悪性高血圧	31	55	2,258	9	1			2,354	264	2,618	2.95	0.28
(%)	(1.3)	(2.3)	(95.9)	(0.4)	(0.0)			(100.0)				
糖尿病性腎症	526	2,071	106,486	339	24	36	7	109,489	10,789	120,278	2.98	0.21
(%)	(0.5)	(1.9)	(97.3)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
SLE腎炎	14	57	1,945	9		2		2,027	200	2,227	2.97	0.26
(%)	(0.7)	(2.8)	(96.0)	(0.4)		(0.1)		(100.0)				
アミロイド腎	3	12	390	2				407	55	462	2.96	0.25
(%)	(0.7)	(2.9)	(95.8)	(0.5)				(100.0)				
痛風腎	9	13	952	3				977	98	1,075	2.97	0.23
(%)	(0.9)	(1.3)	(97.4)	(0.3)				(100.0)				
先天性代謝異常による腎不全	2	6	223			1		232	42	274	2.97	0.31
(%)	(0.9)	(2.6)	(96.1)			(0.4)		(100.0)				
腎・尿路結核		1	179					180	15	195	2.99	0.07
(%)		(0.6)	(99.4)					(100.0)				
腎・尿路結石	4	14	502	3				523	58	581	2.96	0.25
(%)	(0.8)	(2.7)	(96.0)	(0.6)				(100.0)				
腎・尿路腫瘍	6	18	797	1				822	88	910	2.96	0.23
(%)	(0.7)	(2.2)	(97.0)	(0.1)				(100.0)				
閉塞性尿路障害	7	36	603	3				649	81	730	2.93	0.31
(%)	(1.1)	(5.5)	(92.9)	(0.5)				(100.0)				
骨髄腫	2	31	224					257	34	291	2.86	0.37
(%)	(0.8)	(12.1)	(87.2)					(100.0)				
腎形成不全	5	7	542	2				556	104	660	2.97	0.23
(%)	(0.9)	(1.3)	(97.5)	(0.4)				(100.0)				
不明	226	777	25,969	93	10	15	2	27,092	2,805	29,897	2.96	0.27
(%)	(0.8)	(2.9)	(95.9)	(0.3)	(0.0)	(0.1)	(0.0)	(100.0)				
再導入	19	45	1,868	14	1	2	1	1,950	233	2,183	2.97	0.30
(%)	(1.0)	(2.3)	(95.8)	(0.7)	(0.1)	(0.1)	(0.1)	(100.0)				
その他	63	223	6,897	35	2	4	2	7,226	818	8,044	2.96	0.28
(%)	(0.9)	(3.1)	(95.4)	(0.5)	(0.0)	(0.1)	(0.0)	(100.0)				
合計	1,853	5,989	274,889	998	96	139	29	283,993	29,203	313,196	2.97	0.24
(%)	(0.7)	(2.1)	(96.8)	(0.4)	(0.0)	(0.0)	(0.0)	(100.0)				
記載なし	1		13	1				15	6	21	2.93	0.59
(%)	(6.7)		(86.7)	(6.7)				(100.0)				
総計	1,854	5,989	274,902	999	96	139	29	284,008	29,209	313,217	2.97	0.24
(%)	(0.7)	(2.1)	(96.8)	(0.4)	(0.0)	(0.0)	(0.0)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査