

表 2256 体重減少量(kg) 原疾患別（施設血液透析患者 透析回数週3回 透析歴2年以上）

原疾患	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	204	2,342	13,321	23,250	11,231	2,450	450	81	24	44	53,397	1,390	54,787	2.44	0.98
(%)	(0.4)	(4.4)	(24.9)	(43.5)	(21.0)	(4.6)	(0.8)	(0.2)	(0.0)	(0.1)	(100.0)				
慢性腎盂腎炎	14	119	473	707	293	59	9	5	1	1	1,681	34	1,715	2.30	1.01
(%)	(0.8)	(7.1)	(28.1)	(42.1)	(17.4)	(3.5)	(0.5)	(0.3)	(0.1)	(0.1)	(100.0)				
急速進行性糸球体腎炎	6	66	338	502	255	47	5	1		2	1,222	35	1,257	2.37	1.06
(%)	(0.5)	(5.4)	(27.7)	(41.1)	(20.9)	(3.8)	(0.4)	(0.1)		(0.2)	(100.0)				
妊娠腎/妊娠中毒症		42	255	451	139	16	4	1	1		909	22	931	2.27	0.81
(%)		(4.6)	(28.1)	(49.6)	(15.3)	(1.8)	(0.4)	(0.1)	(0.1)		(100.0)				
その他の分類不能の腎炎	6	38	188	297	148	46	3	2			728	9	737	2.42	1.01
(%)	(0.8)	(5.2)	(25.8)	(40.8)	(20.3)	(6.3)	(0.4)	(0.3)			(100.0)				
多発性嚢胞腎	19	285	1,733	2,926	1,280	194	39	1	1	7	6,485	153	6,638	2.38	0.94
(%)	(0.3)	(4.4)	(26.7)	(45.1)	(19.7)	(3.0)	(0.6)	(0.0)	(0.0)	(0.1)	(100.0)				
腎硬化症	65	811	4,032	6,293	2,509	650	127	25	9	10	14,531	351	14,882	2.35	0.99
(%)	(0.4)	(5.6)	(27.7)	(43.3)	(17.3)	(4.5)	(0.9)	(0.2)	(0.1)	(0.1)	(100.0)				
悪性高血圧	9	47	279	537	367	133	30	7	1	1	1,411	40	1,451	2.71	1.12
(%)	(0.6)	(3.3)	(19.8)	(38.1)	(26.0)	(9.4)	(2.1)	(0.5)	(0.1)	(0.1)	(100.0)				
糖尿病性腎症	256	2,896	13,229	25,821	15,870	4,651	971	200	49	50	63,993	1,899	65,892	2.60	1.07
(%)	(0.4)	(4.5)	(20.7)	(40.3)	(24.8)	(7.3)	(1.5)	(0.3)	(0.1)	(0.1)	(100.0)				
SLE腎炎	3	67	340	507	177	42	5			1	1,142	26	1,168	2.26	0.93
(%)	(0.3)	(5.9)	(29.8)	(44.4)	(15.5)	(3.7)	(0.4)			(0.1)	(100.0)				
アミロイド腎	4	13	67	88	29	8					209	7	216	2.12	1.03
(%)	(1.9)	(6.2)	(32.1)	(42.1)	(13.9)	(3.8)					(100.0)				
痛風腎	5	28	145	273	139	35	10	1			636	16	652	2.50	1.01
(%)	(0.8)	(4.4)	(22.8)	(42.9)	(21.9)	(5.5)	(1.6)	(0.2)			(100.0)				
先天性代謝異常による腎不全	2	11	35	48	35	15	2				148	5	153	2.50	1.14
(%)	(1.4)	(7.4)	(23.6)	(32.4)	(23.6)	(10.1)	(1.4)				(100.0)				
腎・尿路結核		8	45	52	12	2	1				120	3	123	2.09	0.83
(%)		(6.7)	(37.5)	(43.3)	(10.0)	(1.7)	(0.8)				(100.0)				
腎・尿路結石	3	22	84	134	60	7	2	1			313	6	319	2.27	0.96
(%)	(1.0)	(7.0)	(26.8)	(42.8)	(19.2)	(2.2)	(0.6)	(0.3)			(100.0)				
腎・尿路腫瘍		20	116	198	97	17	2			1	451	8	459	2.44	0.97
(%)		(4.4)	(25.7)	(43.9)	(21.5)	(3.8)	(0.4)			(0.2)	(100.0)				
閉塞性尿路障害	7	21	115	124	82	18	3	2			372	5	377	2.34	1.10
(%)	(1.9)	(5.6)	(30.9)	(33.3)	(22.0)	(4.8)	(0.8)	(0.5)			(100.0)				
骨髄腫	3	10	25	31	11	3					83	4	87	2.05	1.10
(%)	(3.6)	(12.0)	(30.1)	(37.3)	(13.3)	(3.6)					(100.0)				
腎形成不全		13	73	142	81	20	9	1			339	5	344	2.58	1.02
(%)		(3.8)	(21.5)	(41.9)	(23.9)	(5.9)	(2.7)	(0.3)			(100.0)				
不明	94	957	3,967	6,173	2,796	634	158	33	6	11	14,829	464	15,293	2.36	1.03
(%)	(0.6)	(6.5)	(26.8)	(41.6)	(18.9)	(4.3)	(1.1)	(0.2)	(0.0)	(0.1)	(100.0)				
再導入	4	24	240	458	286	62	9	2	1	1	1,087	21	1,108	2.59	0.94
(%)	(0.4)	(2.2)	(22.1)	(42.1)	(26.3)	(5.7)	(0.8)	(0.2)	(0.1)	(0.1)	(100.0)				
その他	35	267	1,013	1,702	755	190	48	5	1	3	4,019	133	4,152	2.37	1.04
(%)	(0.9)	(6.6)	(25.2)	(42.3)	(18.8)	(4.7)	(1.2)	(0.1)	(0.0)	(0.1)	(100.0)				
合計	739	8,107	40,113	70,714	36,652	9,299	1,887	368	94	132	168,105	4,636	172,741	2.48	1.03
(%)	(0.4)	(4.8)	(23.9)	(42.1)	(21.8)	(5.5)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
記載なし		1	1	1							3		3	1.37	1.19
(%)		(33.3)	(33.3)	(33.3)							(100.0)				
総計	739	8,108	40,114	70,715	36,652	9,299	1,887	368	94	132	168,108	4,636	172,744	2.48	1.03
(%)	(0.4)	(4.8)	(23.9)	(42.1)	(21.8)	(5.5)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy
2015年調査