

表 3116 総コレステロール濃度(mg/dL) 併用の状況別（透析患者全体）

併用の状況	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	8,352	24,359	45,008	52,802	43,512	27,915	14,372	6,523	2,646	1,709	227,198	76,965	304,163	155.92	35.67
(%)	(3.7)	(10.7)	(19.8)	(23.2)	(19.2)	(12.3)	(6.3)	(2.9)	(1.2)	(0.8)	(100.0)				
PD単独	32	134	338	594	714	630	469	287	144	135	3,477	3,627	7,104	181.04	41.44
(%)	(0.9)	(3.9)	(9.7)	(17.1)	(20.5)	(18.1)	(13.5)	(8.3)	(4.1)	(3.9)	(100.0)				
HD(F)単独.PDカテあり	3	17	22	26	31	12	5	4	1		121	61	182	153.97	33.42
(%)	(2.5)	(14.0)	(18.2)	(21.5)	(25.6)	(9.9)	(4.1)	(3.3)	(0.8)		(100.0)				
PD.HD併用週1 HD(F)	17	66	137	198	214	156	95	39	26	13	961	509	1,470	166.98	36.80
(%)	(1.8)	(6.9)	(14.3)	(20.6)	(22.3)	(16.2)	(9.9)	(4.1)	(2.7)	(1.4)	(100.0)				
PD.HD併用週2 HD(F)	2	7	13	29	27	24	12	12	4	3	133	39	172	174.32	37.24
(%)	(1.5)	(5.3)	(9.8)	(21.8)	(20.3)	(18.0)	(9.0)	(9.0)	(3.0)	(2.3)	(100.0)				
PD.HD併用週3 HD(F)	1	1	7	3	2	6	1	1			22	7	29	159.00	35.34
(%)	(4.5)	(4.5)	(31.8)	(13.6)	(9.1)	(27.3)	(4.5)	(4.5)			(100.0)				
PD.HD併用週4 HD(F)															
(%)															
PD.HD併用その他	2	2	5	13	14	6	4	2			48	23	71	161.98	33.26
(%)	(4.2)	(4.2)	(10.4)	(27.1)	(29.2)	(12.5)	(8.3)	(4.2)			(100.0)				
合計	8,409	24,586	45,530	53,665	44,514	28,749	14,958	6,868	2,821	1,860	231,960	81,231	313,191	156.35	35.90
(%)	(3.6)	(10.6)	(19.6)	(23.1)	(19.2)	(12.4)	(6.4)	(3.0)	(1.2)	(0.8)	(100.0)				
不明			1	2	1	1	2	2			9	16	25	184.33	38.01
(%)			(11.1)	(22.2)	(11.1)	(11.1)	(22.2)	(22.2)			(100.0)				
記載なし												1	1		
(%)															
総計	8,409	24,586	45,531	53,667	44,515	28,750	14,960	6,870	2,821	1,860	231,969	81,248	313,217	156.35	35.90
(%)	(3.6)	(10.6)	(19.6)	(23.1)	(19.2)	(12.4)	(6.4)	(3.0)	(1.2)	(0.8)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査