

表 3398 一日透析液使用量(L/日) 原疾患別(腹膜透析患者)

原疾患	2未満	2～	4～	6～	8～	10～	12～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	35	87	334	671	522	138	63	1,850	980	2,830	6.87	2.38
(%)	(1.9)	(4.7)	(18.1)	(36.3)	(28.2)	(7.5)	(3.4)	(100.0)				
慢性腎盂腎炎	3	5	13	26	15	3	4	69	28	97	6.61	2.85
(%)	(4.3)	(7.2)	(18.8)	(37.7)	(21.7)	(4.3)	(5.8)	(100.0)				
急速進行性糸球体腎炎	1	3	3	14	12	6	2	41	25	66	7.38	2.70
(%)	(2.4)	(7.3)	(7.3)	(34.1)	(29.3)	(14.6)	(4.9)	(100.0)				
妊娠腎/妊娠中毒症			5	9	4	2	1	21	6	27	6.95	2.18
(%)			(23.8)	(42.9)	(19.0)	(9.5)	(4.8)	(100.0)				
その他の分類不能の腎炎		3	12	12	10	4	2	43	35	78	7.03	2.71
(%)		(7.0)	(27.9)	(27.9)	(23.3)	(9.3)	(4.7)	(100.0)				
多発性嚢胞腎	9	19	37	68	32	8	2	175	88	263	6.00	2.38
(%)	(5.1)	(10.9)	(21.1)	(38.9)	(18.3)	(4.6)	(1.1)	(100.0)				
腎硬化症	17	84	152	258	157	43	12	723	372	1,095	6.27	2.37
(%)	(2.4)	(11.6)	(21.0)	(35.7)	(21.7)	(5.9)	(1.7)	(100.0)				
悪性高血圧	7	6	15	17	18	1	2	66	41	107	5.88	2.65
(%)	(10.6)	(9.1)	(22.7)	(25.8)	(27.3)	(1.5)	(3.0)	(100.0)				
糖尿病性腎症	39	87	334	645	422	126	43	1,696	1,103	2,799	6.74	2.44
(%)	(2.3)	(5.1)	(19.7)	(38.0)	(24.9)	(7.4)	(2.5)	(100.0)				
SLE腎炎	3	1	6	7	5	4		26	20	46	6.16	3.08
(%)	(11.5)	(3.8)	(23.1)	(26.9)	(19.2)	(15.4)		(100.0)				
アミロイド腎		2	2	2	1			7	8	15	5.16	1.80
(%)		(28.6)	(28.6)	(28.6)	(14.3)			(100.0)				
痛風腎			4	6	8	2	3	23	9	32	8.01	2.37
(%)			(17.4)	(26.1)	(34.8)	(8.7)	(13.0)	(100.0)				
先天性代謝異常による腎不全	1		3	4	4		1	13	4	17	6.78	3.21
(%)	(7.7)		(23.1)	(30.8)	(30.8)		(7.7)	(100.0)				
腎・尿路結核		1		1				2		2	4.75	1.77
(%)		(50.0)		(50.0)				(100.0)				
腎・尿路結石	1	1	2	7	4	1	1	17	4	21	6.72	2.58
(%)	(5.9)	(5.9)	(11.8)	(41.2)	(23.5)	(5.9)	(5.9)	(100.0)				
腎・尿路腫瘍			1	2	4	1		8	6	14	7.66	1.58
(%)			(12.5)	(25.0)	(50.0)	(12.5)		(100.0)				
閉塞性尿路障害		3	4	4	6	1		18	3	21	6.41	2.11
(%)		(16.7)	(22.2)	(22.2)	(33.3)	(5.6)		(100.0)				
骨髄腫			2	1	2			5	2	7	6.70	2.17
(%)			(40.0)	(20.0)	(40.0)			(100.0)				
腎形成不全	4	8	9	6	11	1	9	48	24	72	7.68	5.54
(%)	(8.3)	(16.7)	(18.8)	(12.5)	(22.9)	(2.1)	(18.8)	(100.0)				
不明	13	51	115	173	132	26	11	521	321	842	6.38	2.47
(%)	(2.5)	(9.8)	(22.1)	(33.2)	(25.3)	(5.0)	(2.1)	(100.0)				
再導入	1	1	2	12	10	2	3	31	24	55	8.03	4.52
(%)	(3.2)	(3.2)	(6.5)	(38.7)	(32.3)	(6.5)	(9.7)	(100.0)				
その他	8	11	36	59	28	9	16	167	96	263	7.00	3.91
(%)	(4.8)	(6.6)	(21.6)	(35.3)	(16.8)	(5.4)	(9.6)	(100.0)				
合計	142	373	1,091	2,004	1,407	378	175	5,570	3,199	8,769	6.69	2.56
(%)	(2.5)	(6.7)	(19.6)	(36.0)	(25.3)	(6.8)	(3.1)	(100.0)				
記載なし												
(%)												
総計	142	373	1,091	2,004	1,407	378	175	5,570	3,199	8,769	6.69	2.56
(%)	(2.5)	(6.7)	(19.6)	(36.0)	(25.3)	(6.8)	(3.1)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査