

表 3766 腹膜炎罹患回数(回/年) 透析歴(年)別 (腹膜透析患者)

透析歴	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2年未満	1,974	229	55	11	5	2		1			2,277	7	1,396	3,680
(%)	(86.7)	(10.1)	(2.4)	(0.5)	(0.2)	(0.1)		(0.0)			(100.0)			
2年～	1,604	231	40	15	2	1	1	1		1	1,896	6	1,200	3,102
(%)	(84.6)	(12.2)	(2.1)	(0.8)	(0.1)	(0.1)	(0.1)	(0.1)		(0.1)	(100.0)			
5年～	797	86	18	2	2	1			1		907	2	611	1,520
(%)	(87.9)	(9.5)	(2.0)	(0.2)	(0.2)	(0.1)			(0.1)		(100.0)			
10年～	172	18	4	1							195		119	314
(%)	(88.2)	(9.2)	(2.1)	(0.5)							(100.0)			
15年～	46	6									52		35	87
(%)	(88.5)	(11.5)									(100.0)			
20年～	14	2									16		10	26
(%)	(87.5)	(12.5)									(100.0)			
25年～	5	3									8		4	12
(%)	(62.5)	(37.5)									(100.0)			
30年～	3										3		3	6
(%)	(100.0)										(100.0)			
35年～	1			1							2			2
(%)	(50.0)			(50.0)							(100.0)			
40年～	1										1		1	2
(%)	(100.0)										(100.0)			
合計	4,617	575	117	30	9	4	1	2	1	1	5,357	15	3,379	8,751
(%)	(86.2)	(10.7)	(2.2)	(0.6)	(0.2)	(0.1)	(0.0)	(0.0)	(0.0)	(0.0)	(100.0)			
不明	3	1									4		14	18
(%)	(75.0)	(25.0)									(100.0)			
記載なし														
(%)														
総計	4,620	576	117	30	9	4	1	2	1	1	5,361	15	3,393	8,769
(%)	(86.2)	(10.7)	(2.2)	(0.6)	(0.2)	(0.1)	(0.0)	(0.0)	(0.0)	(0.0)	(100.0)			
平均	2.99	3.04	2.51	3.43	2.56	3.00	2.00	1.50	5.00	2.00	2.98	2.00	3.02	2.99
標準偏差	3.56	3.61	2.59	6.32	2.83	3.16		0.71			3.57	2.00	3.54	3.55

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査