

表 148 2016年導入患者死亡原因分類(年齢別、原疾患別)原疾患 = 糖尿病性腎症

死因	5歳未満	5歳~	10歳~	15歳~	20歳~	25歳~	30歳~	35歳~	40歳~	45歳~	50歳~	55歳~	60歳~	65歳~	70歳~	75歳~	80歳~	85歳~	90歳~	95歳~	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)																					151			151	77.10	9.92
脳血管障害 (%)																					(23.7)			(23.7)		
感染症 (%)																					52			52	70.83	11.52
消化管出血 (%)																					(8.2)			(8.2)		
悪性腫瘍 (%)																					153			153	75.50	10.38
悪液質/尿毒症 (%)																					(24.0)			(24.0)		
心筋梗塞 (%)																					12			12	78.50	8.52
カリウム中毒/頻死 (%)																					(1.9)			(1.9)		
肝硬変症 (%)																					55			55	71.67	8.87
自殺/拒否 (%)																					(8.6)			(8.6)		
腸閉塞 (%)																					27			27	79.22	10.59
肺血栓/肺塞栓 (%)																					(4.2)			(4.2)		
災害・事故死 (%)																					29			29	77.52	7.51
その他 (%)																					(4.5)			(4.5)		
不明 (%)																					11			11	69.36	10.30
合計 (%)																					(1.7)			(1.7)		
記載なし																					6			6	69.67	11.52
総計																					(0.9)			(0.9)		

数値下のかっこ内は列方向の合計に対する%です。

##A083##

(C)Japanese Society for Dialysis Therapy

2016年調査