

表 228 2016年末患者、治療方法(新分類)、原疾患別

原疾患	施設血液透析	Off-line HDF	On-line HDF	Push/Pull HDF	ハイオフィルトレーション	IHDF	血液濾過	血液吸着透析(リクセル等使用)	在宅血液透析	腹膜透析(CAPD)	腹膜透析(APD)	腹膜透析(CCPD)	合計	記載なし	総計
慢性系球体腎炎 (%)	63,328 (69.0)	1,804 (2.0)	19,783 (21.6)	17 (0.0)	90 (0.1)	2,825 (3.1)	9 (0.0)	952 (1.0)	309 (0.3)	1,553 (1.7)	760 (0.8)	353 (0.4)	91,783 (100.0)		91,783
慢性腎孟腎炎 (%)	2,083 (71.9)	50 (1.7)	550 (19.0)	1 (0.0)	5 (0.2)	81 (2.8)		19 (0.7)	10 (0.3)	56 (1.9)	30 (1.0)	11 (0.4)	2,896 (100.0)		2,896
急速進行性系球体腎炎 (%)	1,934 (76.2)	33 (1.3)	428 (16.9)	1 (0.0)	1 (0.0)	68 (2.7)		5 (0.2)	7 (0.3)	36 (1.4)	17 (0.7)	8 (0.3)	2,538 (100.0)		2,538
妊娠腎/妊娠中毒症 (%)	926 (62.4)	53 (3.6)	398 (26.8)	1 (0.1)	2 (0.1)	39 (2.6)		38 (2.6)	2 (0.1)	16 (1.1)	5 (0.3)	3 (0.2)	1,483 (100.0)		1,483
その他の分類不能の腎炎 (%)	984 (67.7)	23 (1.6)	301 (20.7)		1 (0.1)	52 (3.6)		11 (0.8)	7 (0.5)	33 (2.3)	18 (1.2)	24 (1.7)	1,454 (100.0)		1,454
多発性囊胞腎 (%)	8,319 (72.5)	153 (1.3)	2,264 (19.7)	2 (0.0)	8 (0.1)	380 (3.3)		44 (0.4)	29 (0.3)	133 (1.2)	108 (0.9)	31 (0.3)	11,471 (100.0)		11,471
腎硬化症 (%)	24,053 (76.0)	301 (1.0)	4,966 (15.7)	8 (0.0)	23 (0.1)	1,145 (3.6)		28 (0.1)	20 (0.1)	664 (2.1)	337 (1.1)	105 (0.3)	31,650 (100.0)		31,650
悪性高血圧 (%)	1,985 (73.0)	27 (1.0)	490 (18.0)			98 (3.6)		10 (0.4)	5 (0.2)	24 (2.4)	16 (0.9)	2,721 (0.6)	1,454 (100.0)		2,721
糖尿病性腎症 (%)	93,965 (76.0)	1,497 (1.2)	20,691 (16.7)	27 (0.0)	84 (0.1)	4,469 (3.6)	9 (0.0)	75 (0.1)	94 (0.1)	1,602 (1.3)	857 (0.7)	287 (0.2)	123,657 (100.0)		123,657
SLE腎炎 (%)	1,554 (69.9)	52 (2.3)	475 (21.4)		3 (0.1)	57 (2.6)	1 (0.0)	16 (0.7)	9 (0.4)	31 (1.4)	17 (0.8)	7 (0.3)	2,222 (100.0)		2,222
アミロイド腎 (%)	312 (71.2)	12 (2.7)	80 (18.3)			16 (3.7)			2 (0.5)	9 (2.1)	7 (1.6)	438 (100.0)		438	
痛風腎 (%)	776 (73.5)	17 (1.6)	202 (19.1)		2 (0.2)	26 (2.5)		9 (0.9)	1 (0.1)	13 (1.2)	6 (0.6)	4 (0.4)	1,056 (100.0)		1,056
先天性代謝異常による腎不全 (%)	185 (65.8)	5 (1.8)	65 (23.1)			7 (2.5)			2 (0.7)	7 (2.5)	5 (1.8)	5 (1.8)	281 (100.0)		281
腎・尿路結核 (%)	116 (69.0)		39 (23.2)			7 (4.2)		4 (2.4)	1 (0.6)		1 (0.6)		168 (100.0)		168
腎・尿路結石 (%)	431 (74.6)	8 (1.4)	92 (15.9)	1 (0.2)		22 (3.8)		2 (0.3)	1 (0.2)	15 (2.6)	3 (0.5)	3 (0.5)	578 (100.0)		578
腎・尿路腫瘍 (%)	728 (75.9)	15 (1.6)	172 (17.9)	1 (0.1)		30 (3.1)		1 (0.1)		5 (0.5)	6 (0.6)	1 (0.1)	959 (100.0)		959
閉塞性尿路障害 (%)	550 (74.0)	7 (0.9)	134 (18.0)		1 (0.1)	25 (3.4)		5 (0.7)	1 (0.1)	10 (1.3)	7 (0.9)	3 (0.4)	743 (100.0)		743
骨髄腫 (%)	272 (84.2)	8 (2.5)	29 (9.0)		1 (0.3)	5 (1.5)			1 (0.3)	4 (1.2)	3 (0.9)	323 (100.0)		323	
腎形成不全 (%)	407 (61.3)	15 (2.3)	148 (22.3)		2 (0.3)	21 (3.2)		5 (0.8)	1 (0.2)	14 (2.1)	37 (5.6)	14 (2.1)	664 (100.0)		664
不明 (%)	23,184 (73.8)	398 (1.3)	5,732 (18.2)	6 (0.0)	21 (0.1)	1,028 (3.3)		83 (0.3)	71 (0.2)	531 (1.7)	270 (0.9)	88 (0.3)	31,412 (100.0)		31,412
再導入 (%)	1,375 (63.2)	54 (2.5)	566 (26.0)		2 (0.1)	66 (3.0)		27 (1.2)	19 (0.9)	31 (1.4)	15 (0.7)	19 (0.9)	2,174 (100.0)		2,174
その他 (%)	6,237 (73.9)	105 (1.2)	1,511 (17.9)	1 (0.0)	6 (0.1)	261 (3.1)		21 (0.2)	25 (0.3)	144 (1.7)	104 (1.2)	23 (0.3)	8,438 (100.0)		8,438
合計 (%)	233,704 (73.2)	4,637 (1.5)	59,116 (18.5)	66 (0.0)	252 (0.1)	10,728 (3.4)	19 (0.0)	1,355 (0.4)	617 (0.2)	4,973 (1.6)	2,637 (0.8)	1,005 (0.3)	319,109 (100.0)		319,109
記載なし (%)															
総計 (%)	233,704 (73.2)	4,637 (1.5)	59,116 (18.5)	66 (0.0)	252 (0.1)	10,728 (3.4)	19 (0.0)	1,355 (0.4)	617 (0.2)	4,973 (1.6)	2,637 (0.8)	1,005 (0.3)	319,109 (100.0)		319,109

数値下の括弧内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

##A0163##

(C)Japanese Society for Dialysis Therapy

2016年調査