

表 302 2016年死亡患者死原因分類(年齢別、原疾患別)原疾患 = 慢性糸球体腎炎

死因	5歳未満	5歳~	10歳~	15歳~	20歳~	25歳~	30歳~	35歳~	40歳~	45歳~	50歳~	55歳~	60歳~	65歳~	70歳~	75歳~	80歳~	85歳~	90歳~	95歳~	合計	不明	記載なし	総計	平均	標準偏差			
心不全 (%)								1	3	16	41	85	211	216	332	407	315	163	34	1,842			1,842	78.13	9.86				
脳血管障害 (%)								1	3	6	8	6	23	30	76	60	69	80	49	15	6	432			432	73.53	11.53		
感染症 (%)									3	4	17	28	73	144	174	252	330	271	114	20	1,431			1,431	78.31	9.49			
消化管出血 (%)											1	4	6	12	15	20	11	12	5	3	89			89	76.02	9.87			
悪性腫瘍 (%)											4	16	31	48	127	111	101	110	95	26	5	680			680	74.14	10.46		
悪液質/尿毒症 (%)											3	1	9	15	18	50	74	66	29	10	275			275	81.22	8.54			
心筋梗塞 (%)												3.1	(0.5)	(2.5)	(1.8)	(2.0)	(4.2)	(5.2)	(6.0)	(6.0)	(9.3)	(4.1)			(4.1)				
カリウム中毒/頻死 (%)												1	3	4	8	26	34	41	40	47	24	7	3	238			238	74.33	10.03
肝硬変症 (%)												2	5	9	13	18	27	32	34	14	5	1	161			161	73.89	10.87	
自殺/拒否 (%)												1	2	2	4	6	11	10	13	7	6	62			62	71.13	10.77		
腸閉塞 (%)												1	2	2	2	5	6	7	8	13	1	2	49			49	76.39	13.28	
肺血栓/肺栓栓 (%)												1	1	1	3	15	18	14	14	6	3	76			76	74.36	9.48		
災害・事故死 (%)												1	2	1	3	2	5	3	1	1	19			19	70.84	12.92			
その他 (%)												1	2	2	1	7	5	10	8	5	2	44			44	73.20	12.27		
不明 (%)												4	4	8	17	29	59	72	119	162	138	62	15	689			689	78.76	10.06
合計 (%)								1	3	2	7	27	54	98	199	365	824	895	1,193	1,426	1,099	483	107	6,783			6,783	77.00	10.27
記載なし																													
総計								1	3	2	7	27	54	98	199	365	824	895	1,193	1,426	1,099	483	107	6,783			6,783	77.00	10.27

数値下のかっこ内は列方向の合計に対する%です。

##A0237##

(C)Japanese Society for Dialysis Therapy

2016年調査